

PERSPECTIVES ON ANIMAL BIOSCIENCES

Shelter and shade for grazing sheep: implications for animal welfare and production and for landscape health

David G. Masters^{A,B,*} , Dominique Blache^A, Amy L. Lockwood^C , Shane K. Maloney^D, Hayley C. Norman^B , Gordon Refshauge^E and Serina N. Hancock^C

For full list of author affiliations and declarations see end of paper

***Correspondence to:**

David G. Masters
School of Agriculture and Environment,
Faculty of Science, The University of
Western Australia, 35 Stirling Highway,
Crawley, WA 6009, Australia
Email: david.masters@uwa.edu.au

Handling Editor:

Alan Tilbrook

Received: 12 June 2022
Accepted: 4 January 2023
Published: 30 January 2023

Cite this:

Masters DG et al. (2023)
Animal Production Science, **63**(7), 623–644.
doi:[10.1071/AN22225](https://doi.org/10.1071/AN22225)

© 2023 The Author(s) (or their
employer(s)). Published by
CSIRO Publishing.
This is an open access article distributed
under the Creative Commons Attribution-
NonCommercial-NoDerivatives 4.0
International License (CC BY-NC-ND).

OPEN ACCESS

ABSTRACT

Shade and shelter may provide protection from cold and heat stress, a source of feed during prolonged or seasonal drought, specific essential nutrients, increased pasture and crop production and improved landscape health. Cold stress contributes to the average of 8% (single) and 24% (twin) of lambs that die within 3 days of birth in Australia and the estimated 0.7% of the Australian flock that die post-shearing during extreme or unseasonal weather. Shelter has resulted in an average reduction in mortality of 17.5% for twin-born lambs and 7% for single-born lambs according to Australian studies and decreases the susceptibility of ewes to metabolic disease and possibly dystocia. Because many of the published studies are from research areas where cold stress is expected, they are not indicative of industry-wide responses, a research priority is to determine the probability of lamb and ewe deaths from cold stress across different sheep production areas. Although shelter may improve lamb survival, ewes do not always choose to lamb in a sheltered location. For this reason, there is a requirement for research into the voluntary use of shelter in commercial-sized paddocks and the role that nutritive value of shelter plays in attracting and holding ewes to shelter, and to their lambs. Heat stress may also result in lamb deaths and influences feed conversion efficiency, appetite, reproduction, wool growth and disease susceptibility. The consequences of heat stress may go unnoticed over a yearly production cycle, although there is some evidence that shade may increase weaning rates and feed intake of grazing sheep. There are ancillary benefits from shade and shelter. Trees may improve crop production through reducing wind damage and evapotranspiration and provide timber. Shrubs provide feed during the summer–autumn feed gap or drought, are useful for the management of land degradation and provide habitat for native fauna. It is clear that shade and shelter in the correct locations provide a range of benefits to livestock and the landscape; nevertheless, adoption appears low. Research that focuses on defining the benefits on a farm or landscape scale is required to support extension programs.

Keywords: chill index, cold stress, drought feeding, heat stress, hypothermia, lamb survival, nutrition, shrubs.

Introduction

Shelter for grazing sheep provides protection from cold, wet and windy conditions or shade from excessive summer heat. The Australian Animal Welfare Standards and Guidelines for Sheep state the guideline ‘that sheep and lambs should be provided with adequate shelter. In the absence of natural protection, consideration should be given to the provision of shade, windbreaks or sheds’ (p. 11) (Animal Health Australia (AHA) 2016). Most shelter research has focused on reducing cold stress on lambs and lambing ewes. This is justified as an average of 8% (single) and 24% (twin) of all lambs born in Australia die within 3 days of birth (80% of total lamb deaths; Hinch and Brien 2014). High lamb mortalities have been reported in the UK (10–25%) and New Zealand (Mellor and Stafford 2004; Celi and Bush 2010), also with a high proportion of deaths in the first 48 h (Dwyer 2008). Similar mortalities would be expected in all cool or temperate

climates where sheep are raised extensively. The extent to which hypothermia is responsible for these lamb deaths is uncertain as it may be part of a complex where starvation, birth injury and low birth weight increase susceptibility to cold. Cold stress during lambing may also contribute to ewe deaths either directly or through interactions with dystocia. There are few reliable data on the deaths of ewes during lambing although one study indicates ewe mortality of 2.5–3%, with 29–41% of these attributable to dystocia (McQuillan *et al.* 2021).

There have been many media articles, anecdotal reports and a few historical scientific studies (Geytenbeek 1962; Hutchinson 1968; Buckman 1982) on sheep deaths during unseasonal storms. Sometimes, but not always, these are associated with very low temperatures. Under such circumstances, significant benefits of shelter may be expected. A survey of sheep losses following the passage of a rain bearing depression during summer in south-Western Australia indicated that 27% of recently shorn sheep died during or shortly after the rain. Most sheep died from hypothermia, even though air temperatures were not low (13.3–21.7°C); conversely, sheep housed in sheds during the storm had close to zero mortality (Buckman 1982).

Heat stress may influence survival and production of both ewes and lambs. The consequences are probably underestimated, with few reports on the production responses from providing shade under commercial conditions. Arnold and Morgan (1975) reported 26–29% mortality from heat stroke in lambs born in summer. This may be a problem for the very few sheep farms in Australia that lamb during hot months (Crocker *et al.* 2009), but is unlikely to be a serious industry problem as few farms in the major sheep production regions in Australia lamb during high temperatures (Hacker 2010). More importantly, under controlled conditions, heat stress has been shown to influence feed conversion efficiency, appetite, reproduction, wool growth and susceptibility to disease (Thwaites 1967, 1971; Sawyer 1979; Morrison 1983; Silanikove 2000; Marai *et al.* 2007; Phillips 2016; Moslemipur and Golzar-Adabi 2017). Under grazing conditions, similar changes may go unnoticed over a yearly production cycle.

Given the impact of heat and cold stress on lambing ewes and newborn lambs in winter and summer and on adult sheep during unseasonal rain and wind, benefits of shade and shelter would be expected. These benefits may have both economic and welfare implications. The use of shelter during lambing has been well researched and reviewed (Radcliffe 1983; Gregory 1995; Hawke and Dodd 2003; Pollard 2006), but there has been less research into the commercial production benefits of shelter and shade outside the lambing period, including the benefits on quantity and quality of the feedbase and on landscape health. The purpose of this review is to summarise research that has been conducted, identify potential production and welfare benefits that have not been evaluated, provide some preliminary guidelines on the use of shelter and shade and,

present priorities for future research and extension. The focus is on the Australian sheep industry, but relevant information from elsewhere is also presented.

Changes in production and reproduction resulting from the use of shelter and shade

Metabolic consequences of cold and heat stress

The metabolic response of a homeothermic animal, such as a sheep, to different environmental conditions can be expressed as a heat balance equation. In heat balance, heat inputs to the body are balanced by heat loss from the body. The generation of metabolic heat forms the central component of the equation, as follows:

$$\text{MHP} \pm \text{Cond} \pm \text{Conv} \pm \text{Rad} \pm \text{Evap} \pm \text{Work} \pm \text{Storage} = 0$$

where MHP is metabolic heat production, and this term is always positive; Cond is heat exchange by conduction and depends on the temperature and thermal conductivity of anything that the body touches; Conv is heat exchange by convection and depends on the air temperature and wind speed; Rad is heat exchange by radiation and depends on the difference between emitted radiation, that depends on the fourth power of the surface temperature of the body, and received radiation, that depends on the fourth power of every surface that the body can ‘see’ and the emissivity (reflectivity) of the body to that received radiation; Evap is heat exchange by evaporation (or condensation), and in nearly all conditions this term is negative and can be modified by panting and sweating; Work is work against the environment, and for a sheep that is not moving, the term = 0; and Storage is heat storage in the body; any imbalance in the sum of the other terms results in either gain of heat to, or loss of heat from, the body. The change in body temperature then is a function of the heat capacity of the body, on average 3.5 J/g.°C.

Heat exchange by the routes above (summarised in Fig. 1) depends on the surface temperature of the body, which in turn depends on blood flow to the periphery. Metabolic heat is generated mainly in internal organs of the body and the heat produced by those organs is carried away from the organs and to the skin and respiratory tract by the blood. Thus, an increase in peripheral blood flow will result in an increase in skin temperature.

There is a range in environmental temperature, known as the thermoneutral zone (TNZ), within which a sheep can maintain body temperature by adjusting blood flow to the periphery, and without resort to extra metabolic heat production or evaporative heat loss (Fig. 1). The TNZ is framed by the lower critical temperature (LCT) at its lower end, and by the upper critical temperature (UCT) at its upper end. Below the LCT, a sheep will increase metabolic




Fig. 1. Principles of thermoregulation in the sheep (adapted from both Porter and Gates (1969) and Mitchell *et al.* (2018)).

heat production to maintain heat balance, while above the UCT it will increase evaporative heat loss (see Fig. 1). The LCT is defined by a single environmental variable, air temperature, while heat balance is affected by many other factors. Thus, the LCT varies widely depending partly on animal factors (fleece length, feed intake, subcutaneous insulation, and posture) as well as environmental factors (wind speed, rainfall, and solar radiation). Anything that increases heat loss (removal of fleece, more wind, more rain) will increase the LCT. Gregory (1995) reported that the LCT for an adult sheep ranges from as high as 28°C when the sheep are shorn, wet, fed at maintenance and exposed to wind, to as low as -35°C when they have a full, dry fleece and are not exposed to wind. Similarly, Freer *et al.* (2007) reported that for 5 kg lambs, the LCT ranged from 28°C when the lambs had 8 mm wool, wind speed was 20 km/h and daily rainfall was 10 mm, to 18°C when the lambs had 14 mm of fleece and were under calm, dry conditions.

When the LCT is increased, more energy must be diverted from growth or production to the production of heat for maintenance of body temperature. That energy can be sourced from body stores or from increased feed intake. Using the equations of Freer *et al.* (2007), it can be estimated that the additional energy cost (above thermoneutral maintenance) to maintain body temperature of a wet 5 kg lamb with 8 mm of wool coverage at 5°C and 20 km/h of wind would be 3.0 MJ ME/day. This energy is additional to the maintenance requirement of the lamb and is equivalent to the energy required to support almost 200 g/day of growth in a

thermoneutral environment (Agricultural Research Council 1980). The additional energy requirements would be lowered to 2.6 MJ ME/day if wind speed could be reduced by 10 km/h by using shelter (Fig. 2).

For an adult 50 kg shorn sheep approximately 7.3 MJ ME/day is required for maintenance under thermoneutral conditions (Freer *et al.* 1997). That requirement is increased by 13.9 MJ ME/day by exposure to 5°C,




Fig. 2. Energy required to maintain liveweight of a 5 kg lamb with 8 mm wool cover in a wind of 20 km/h (solid line), 10 km/h (short, dashed line) or 0 km/h (long dashed line). All scenarios following 10 mm rainfall. Derived from equation 1.24 from Freer *et al.* (2007).




Fig. 3. Energy required to maintain liveweight of a 50 kg ewe with (a) 5 mm or (b) 50 mm wool cover in a wind of 20 km/h (solid line), 10 km/h (short, dashed line) or 0 km/h. All scenarios following 10 mm rainfall. Derived from equation 1.24 from Freer *et al.* (2007).

20 km/h of wind and 10 mm of rain (Fig. 3a). If wind speed is reduced to 10 km/h, the extra energy expenditure would be reduced to 12.1 MJ ME/day (Fig. 3a). Insulation with 50 mm of wool under similar weather conditions means that energy requirements increase by only 2.8 MJ ME/day (Fig. 3b).

For both lambs and shorn adults, the generation of enough heat through increased intake would be a challenge. Measured increases in feed intake of 46–78% in shorn sheep exposed to 7–10°C (Elvidge and Coop 1974) are consistent with the predictions of Freer *et al.* (2007). It is unlikely that shorn adults grazing short winter or dry summer pastures would be able to consume enough additional feed to meet the energy requirements of exposure to cold and rain. Subsequently, weight loss would be expected with a reduction in the efficiency of use of available nutrients for growth and production.

Even when sufficient energy is available from body stores or extra intake, there is another limit on the generation of heat. The maximum metabolic rate is known as summit metabolism, and under extreme cold stress the metabolic heat production that is required to achieve heat balance can exceed the summit metabolism (Fig. 1). Under those conditions, heat loss will exceed heat gain, hypothermia will develop, tissue oxygenation will decrease (hypoxia) and/or the partial pressure of oxygen in the blood will decrease (hypoxaemia) followed by death, regardless of feed availability or body stores. There are conflicting conclusions on whether summit metabolism increases following colostrum intake (Alexander 1962; Eales and Small 1980).

There is also an energy cost associated with heat stress, at least in sheep under controlled conditions. Those costs are given as 7% for rapid shallow panting (often called Phase I panting) and 11–25% for deep open-mouth panting (often called Phase II panting; National Research Council 1981). In a commercial production system, the total energy cost of heat stress is more difficult to quantify because temperature, other environmental parameters and animal behaviour that can mitigate stress all vary within the day.

Cold stress, shelter and lamb survival, health and production

In a study on Kangaroo Island between 1959 and 1963, the highest daily rate of lamb mortality was 54% for Corriedales and 91% for Merinos. These high rates occurred on days when wind speed was 24–56 km/h, rainfall ranged from 5.3 to 26 mm/day and average temperature ranged from 8.6°C to 12.3°C (Obst and Day 1968). Mortality was significantly higher when wind speed exceeded 8 km/h and daily rainfall exceeded 0.25–5 mm/day than on days when there was no wind or rain (Obst and Day 1968). Donnelly (1984) analysed data collected near Canberra between 1970 and 1971 and reported that the probability of death from exposure in the first 3 days of life for Merino lambs was up to 40% for singles and 60% for multiples. For crossbred lambs, the reported probabilities were 25% for singles and 40% for multiples. On low-chill days, the mortality of lambs born to ewes in good condition approached zero (Donnelly 1984). Across a range of environments, Dennis (1974) autopsied 4417 lambs from farms in Western Australia between 1963 and 1965 and reported exposure as the cause of death for 2.1% of lambs. More recently, Refshauge *et al.* (2016) autopsied lambs that died within 5 days of birth at sites across Australia between 2008 and 2011 and found that, on average, 5% of lambs died due to exposure. However, death varied among sites and years, with between 0% and 38% of lambs dying from hypothermia.

Low-birthweight lambs (often multiples) are at a higher risk of death than are heavy lambs (Refshauge *et al.* 2016). For survival, summit metabolism must meet or exceed heat loss. Summit metabolism is 71 kJ/kg.h for both single and

twin lambs (Alexander 1962). However, heat loss is related to skin-surface area and summit metabolism per unit of surface area decreases as bodyweight decreases; this means larger, heavier lambs are better able to maintain body temperature than are small lambs, irrespective of birth status. When the cold-stress conditions on a lamb are below the tolerance zone (Fig. 1), summit metabolism is insufficient to balance heat loss, and hypothermia develops. A positive feedback cycle results because summit metabolism decreases as body temperature declines (Alexander 1962).

The benefits of shelter under such situations can be high; in a comprehensive 5-year study involving Merinos in New South Wales, where at least 40% of the lamb mortality was associated with hypothermia (Alexander *et al.* 1980), grass tussocks reduced the mortality in singles from 17% to 9% and in multiples from 51% to 36%. Wind protection alone saved many single lambs during poor weather but was less effective in protecting multiples unless the weather was fine. Low-birthweight multiples were more susceptible to hypothermia than were heavier lambs and prolonged poor weather resulted in hypothermia even in older lambs that had consumed milk. The observation that shelter improves survival has been confirmed in many other studies (McLaughlin *et al.* 1970; Egan *et al.* 1972, 1976; Robertson *et al.* 2011); these are summarised in Table 1.

The extent of the reduction in mortality is weather related, with many studies concluding that only twin lambs benefit from shelter. Twins are usually smaller than singles and more susceptible to high chill. Single-lamb mortality is often dismissed as non-significant (Egan *et al.* 1976; Robertson *et al.* 2011). However, while often not statistically significant, there is a consistent decline in death rates of single lambs (0–13%) as well as twin lambs (11–34%) at high-risk locations (Table 1).

In assessing the benefits of shelter, it is important to consider that many published results focus on locations, climates and events where lambing coincides with periods of high chill index. The chill index, which was derived by Nixon-Smith (1972) and modified by Donnelly (1984), is an indicator of potential heat loss and is calculated from temperature, wind velocity and rainfall. The studies undertaken in high-risk locations have well described the potential consequences but not the probability of those consequences. When weather conditions are mild and the chill index low, deaths due to cold stress are also low (Obst and Day 1968) and lamb survival is not improved by providing wind breaks (Lynch and Alexander 1976). Defining when shelter will be beneficial is not simple but can be partially addressed using the methodology of McCaskill and Clark (2007). They constructed wind roses to coincide with wind directions at times of bad weather during lambing and could quantify the percentage of time when the weather was bad enough to cause mortality. In a more comprehensive study (Broster *et al.* 2012a), the seasonal risk of cold stress was defined over 7-day periods

for 24 weeks at six sheep growing locations in south eastern Australia. Weather conditions including wind, temperature, and rainfall were modelled using the GrassGro decision support tool (Donnelly *et al.* 1997). During the coldest months of the year, the median chill index exceeded 1000 kJ/m².h for nearly 100% of the time at some sites and less than 20% of the time at others. The threshold chill index of 1000 kJ/m².h was based on estimates of the summit metabolism in newborn lambs of 837–1255 kJ/m².h (Alexander 1962). A reduction in wind speed of 25% significantly reduced the median and maximum chill indices (Broster *et al.* 2012a). A key point from this study was that shelter will be far more effective in reducing chill index at windy locations and less effective where chill index is driven by temperature. A minimum starting point to encourage the broad establishment of shelter is to assess the risks at lambing time and across all sheep-growing regions by using methods similar to those of Broster *et al.* (2012a) or McCaskill and Clark (2007), and to use this information to estimate the expected number of lambs that could be saved, and calculate the financial benefit from the establishment of shelter at different locations.

Cold stress, shelter, ewe health, survival and production

Ewes, particularly those with a long fleece, have a very high cold tolerance (Fig. 3b). However, under extreme conditions, cold stress will compromise health and survival often through an interaction with body condition and metabolic disease. A combination of cold or emotional stress and short periods of starvation can induce pregnancy toxemia, particularly if ewes are in poor condition (Reid 1962). Pregnancy toxemia is a metabolic disease that can occur during late pregnancy when high energy requirements are not being met or when feed intake is depressed (Mavrogianni and Brozos 2008). The risk of energy imbalance is enhanced by an increase in energy requirements for the maintenance of body temperature and a possible reduction in digestive efficiency during cold stress (Young 1981). The resulting catabolic state causes high blood ketone concentrations and low blood glucose with increased ewe and fetal mortality. Early observations have described how adverse weather hastened the onset of 'undernutrition syndrome' due to a reduction in feed intake and an increase in energy requirement (Reid 1962). Paradoxically, the reduced intake and stress may be due to movement of the animals to a sheltered area without edible plants (Andrews 1997). The inappetence associated with pregnancy toxemia can further result in hypocalcaemia and/or hypomagnesaemia. These metabolic diseases result from an acute shortage of available calcium or magnesium and can cause mortality of the ewe and fetus and, in a subclinical form, may be associated with uterine inertia and prolonged parturition (Friend *et al.* 2020).

Table 1. Influence of shelter on lamb mortality from experiments conducted in Australia between 1968 and 2011.

Lambing birth type	Breed	Lamb mortality (%)			Shelter type	Paddock size (ha, except when indicated otherwise)	Location	Reference
		Sheltered lambing	Exposed lambing	Shelter benefit				
Single	Merino	9	17	8	Grass hedge	0.4	Armidale, NSW	Alexander et al. (1980)
Single	Merino	6	19	13	Trees/shrubs	0.8–1.5	Hamilton, Vic.	Egan et al. (1972)
Single	Merino	10	12	3	Grass	0.1	Hamilton, Vic.	Egan et al. (1976)
Single	Corriedale	7	13	7	Trees/shrubs	1.5–2.5	Hamilton, Vic.	McLaughlin et al. (1970)
Single	Merino cross	22	25	0	Hedge/shrubs	1	Tarcutta, NSW	Robertson et al. (2011)
Single	Corriedale	8	21	13	Various	Pens	Western Vic.	Watson et al. (1968)
Single	Merino	5	11	6	Grass hedges	<0.4–0.8	Armidale, NSW	Lynch and Alexander (1977)
Average				7				
Twin	Merino	13	27	15	Trees/shrubs	0.8–1.5	Hamilton, Vic.	Egan et al. (1972)
Twin	Merino	19	31	12	Grass	1.5–2.5	Hamilton, Vic.	Egan et al. (1976)
Twin	Corriedale	6	25	19	Trees/shrubs	1.5–2.5	Hamilton, Vic.	McLaughlin et al. (1970)
Twin	Merino	41	68	27	Grass hedges	0.4–0.8	Armidale, NSW	Lynch and Alexander (1977)
Twin	Merino cross			>4.0	Trees/shrubs	1.3	Tarcutta, NSW	Robertson et al. (2011)
Twin		24	58	34	Various	Pens	Western Vic.	Watson et al. (1968)
Twin	Merino	27	12	15	Oats, double row	1	Wooroloo, WA	Glover et al. (2008)
Twin	Merino	27	26	1	Oats, single row	1	Wooroloo, WA	Glover et al. (2008)
Average				17.5				
Multiple	Merino	36	51	15	Grass hedges	0.4	Armidale, NSW	Alexander et al. (1980)
Singles + twins	Merino	21	39	17	Grass hedges + polythene mesh	12	Armidale, NSW	Lynch et al. (1980)

Cold stress can also directly influence survival. In Australia, historical data indicate that the mortality of ewes and lambs post-shearing is 0.7% of all sheep (Hutchinson 1968). Over 75% of these deaths are within 14 days of shearing (Hutchinson 1968). Individual flocks and regions can have a much higher death rate than the average. Geytenbeek (1962) reported mortality within 5 days of shearing of 13.9% for Merinos and 8.2% for Corriedales and Crossbreds from a survey of 17 properties in the Kybybolite region of South Australia. Overall mortality rates of 26–28% have been reported for recently shorn sheep in the south-western

and central wheatbelt in Western Australia under poor weather conditions (Buckman 1982; Glass and Jacob 1992). Glass and Jacob (1992) observed mortality of 90% on one farm where sheep were shorn less than a week before adverse weather. Hence, the risk of mortality after shearing is exacerbated with heavy rain and a cold wind, particularly when sheep are in poor body condition (Geytenbeek 1962; Hutchinson 1966, 1968). Deaths may be reduced to zero with adequate shelter, although high mortalities have been recorded even when paddocks were considered well sheltered (Geytenbeek 1962; Buckman 1982; Glass and Jacob 1992).

This may be due to the positioning of shelter being ineffective against the prevailing winds, insufficient density of shelter at the level of the sheep, shelter being provided too late or extended periods of exposure to heavy rains (Glass and Jacob 1992).

It would be expected that the requirement for energy to be diverted for the maintenance of body temperature during cold stress would mean a depression in growth and production. In Welsh Mountain ewes under cold conditions and controlled feed intake, outdoor exposure led to a 1.5 kg reduction in bodyweight pre-partum, equivalent to a 10% reduction in daily digestible organic-matter intake. The growth rate of wool was unchanged (Robinson *et al.* 1973). Conversely, in Scottish Blackface sheep wool growth was reduced under cold conditions in climate-controlled chambers (Slee and Ryder 2009). The diversion of energy to thermoregulation affects other physiological systems. An artificially induced stress that included simulated rain in a cold environment reduced ovulation rate by 8–42% (Doney *et al.* 1973). Cold stress on the ewe will also influence the development and subsequent survival of the fetus. Lambs from recently shorn ewes or ewes in poor condition are more likely to die from hypothermia than lambs from unshorn, well fed ewes (Alexander *et al.* 1980; Donnelly 1984; Oldham *et al.* 2011), although shearing does not always increase the risk of mortality if shelter is available (Lynch and Alexander 1976). Shearing in mid-pregnancy may also result in increased birthweight, although results are inconsistent (Kenyon *et al.* 2003). There is a well documented relationship among ewe condition, lamb birthweight and lamb survival. Ewes in better condition tend to be more insulin-insensitive, resulting in the partitioning of more glucose to the fetus and in heavier lambs with higher fat reserves that are less likely to die around the time of birth (McNeill *et al.* 1997). Less well understood are the possible effects of thermal stress on gene expression in the ewe and fetus and the potential for epigenetic changes. These are apparent in cultured cells (Sonna *et al.* 2002), with some evidence of changes in living organisms (Abe *et al.* 2018).

There are few reports on the direct impact of shelter on cold-stressed, adult sheep under commercial conditions. In one study, there were fewer shepherding problems (dystocia or mismothering) in ewes with shelter (Pritchard *et al.* 2021); however the number of ewes in this unreplicated study was small. The lack of reports on changes in production are possibly due to an increase in feed intake and compensatory growth under extensive production. Nevertheless, because energy is used for heat production, cold stress would reduce the efficiency of feed use for production and therefore also reduce the sustainable stocking capacity, particularly when grazing early season, slow growing winter pastures.

Heat stress, shelter and ewe and lamb survival, health and production

Controlled environment

Much of the research into the consequences of heat stress on sheep has been under controlled conditions, often in a climate-controlled facility. With sustained heat stress (days rather than hours), feed intake, feed digestibility, liveweight and wool growth are reduced, water intake increased (Thwaites 1967; Marai *et al.* 2007), and early embryo death and returns to service are both increased (Dutt *et al.* 1959; Smith *et al.* 1966; Thwaites 1971). Heat stress disrupts the oestrus cycle, slows placental growth, depresses lamb number, and results in low-birthweight lambs (Sawyer 1979; McCrabb *et al.* 1993; van Wettere *et al.* 2021). Exposure to high temperatures even for short periods prior to mating (32°C on the 12th day of the cycle before breeding) significantly reduces ova fertilisation (Dutt *et al.* 1959; Dutt 1964). In males heat stress is associated with fewer sperm per ejaculate, abnormal sperm and reduced sperm motility (Smith 1971). Short-term heat stress is associated with elevated oxidative stress, increased cortisol and catecholamines and reduced thyroid hormones in plasma (Silanikove 2000; Lowe *et al.* 2002; van Wettere *et al.* 2021). Blood flow changes (Hales 1973), and panting increases the risk of alkalosis (Barnes *et al.* 2004). These changes are associated with altered neurological function with the lowered partial pressure of carbon dioxide in the cerebral blood vessels causing vasoconstriction and reduced perfusion of the brain (Stockman *et al.* 2011). High humidity exacerbates the heat stress that is caused by high ambient temperature because evaporation is not as effective when humidity is high.

Grazing sheep

In sheep flocks grazing in tropical and arid conditions, high temperatures cause lower lamb birth weight and often the death of newborn lambs (Arnold and Morgan 1975; Hopkins *et al.* 1980). The availability of shade can significantly increase the weaning percentage (Stephenson *et al.* 1984). Under extensive grazing in temperate or Mediterranean environments, variation in temperature and humidity across and within days, combined with behavioural thermoregulation, can reduce the likelihood of the sustained heat stress that is observed under controlled or tropical conditions (Brown 1971); nevertheless, reduced embryo survival and fertility, along with increased returns to service, have been associated with high temperatures during or just after mating (Kleemann and Walker 2005). Even short periods of heat stress in grazing ewes can depress appetite and compromise placental growth (McCrabb and Bortolussi 1996). In the field, this heat stress can interact with other environmental and nutritional factors to exacerbate the issue; for example, digestion of the low-quality pastures that are often associated with high environmental temperatures will generate more heat per unit of metabolisable energy (ME)

and therefore contribute more to the overall heat load on the animal (Webster *et al.* 1975). Increased walking for water and feed will increase energy use and heat production. Panting can disrupt the acid–base balance and increase the requirements for mono-valent ions such as sodium, potassium, and chlorine (Silanikove 2000; Barnes *et al.* 2004). In sheep exposed to heat in an indoor facility, the oxidative stress that is associated with heat can be partially alleviated by feeding supranutritional amounts of antioxidants such as vitamin E (Chauhan *et al.* 2014a, 2014b).

The consequences of heat stress in grazing sheep may not therefore be immediately apparent but will affect long-term growth and reproduction. There is sufficient evidence to indicate that the reduction of heat stress should be a priority, particularly during mating and pregnancy.

Because radiation is a major avenue of heat gain on cloud-free days (Fig. 1), benefits of shade to sheep under heat stress would be expected; but benefits have not been conclusively demonstrated. Johnson and colleagues (Johnson 1987, 1991; Sherwin and Johnson 1990; Johnson and Strack 1992) investigated the provision of shade in sheep at maximum ambient temperatures up to 50°C and observed few differences in feeding, drinking or ruminating between sheep that chose to use shade and those that had shade available but chose not to use it. Silanikove (1987) reported that shade reduced the respiration rate and body temperature but had no effect on bodyweight or feed intake. In a later publication, Silanikove (2000) concluded that the incidence of heat stress is highly dependent on shade, feed amount and composition and water availability and quality. Much of the support for the benefit of shade has been based on responses in cattle rather than sheep.

Changes in behaviour resulting from the use of shelter and shade

Cold

How the ewe and lamb choose to interact with shelter and with each other is critical for achieving the benefits that are available. To gain protection from wind and possibly rain, the ewe will need to voluntarily seek shelter during poor weather and during lambing. Most studies on shelter and behaviour during lambing have been in small paddocks, plots, or even sheds (Table 1). There have been few studies on behaviour and voluntary use of shelter at the commercial scale. In small plots, grass rows as far apart as 240 m were used by recently shorn, lambing ewes, but less so by unshorn sheep. Shearing ewes 4–9 days prior to expected lambing increased the use of shelter during both nights and days, either before or after lambing (Alexander *et al.* 1979). Shelter was used more at night and during bad weather by shorn (but not unshorn) ewes (Mottershead *et al.* 1982). Grass shelter was preferred to black plastic mesh, but many ewes preferred to lamb in

isolation, away from shelter (Alexander *et al.* 1979). In contrast, Broster *et al.* (2017) observed that a higher than expected number of ewes lambed close to shelter rows and in New Zealand experiments a high proportion of lambs used shelter on commercial farms (Pollard and Littlejohn 1999). Stevens *et al.* (1981) reported that Merino ewes in moderate-sized paddocks did not seek isolation at lambing and tended to lamb where the flock was grazing or resting at the commencement of parturition. A high proportion of ewes lambed in shelter because this was where the flock was located at the time. Others have reported frequent use of shelter in ewes in moderate-sized paddocks (Lynch *et al.* 1980; Taylor *et al.* 2011). The inconsistent conclusions warrant further investigation to determine whether sheep will seek shelter in large paddocks or whether shelter can be made more attractive, or whether small, specialised lambing paddocks are required.

During the postnatal period, ewe appetite will increase. This too may influence behaviour. Studies on interactions between food preference and social bonds have indicated that ewes will try to maximise energy gain (Dumont *et al.* 1998), and forage availability and quality will influence the distance that ewes will travel away from social groups (Dumont and Boissy 2000). Energy may not be the only nutritional attraction. During late pregnancy and parturition, ewes are susceptible to metabolic disturbances related to calcium, magnesium, sodium and potassium availability such as hypocalcaemia and hypomagnesaemia. The transition period (3 weeks before to several weeks after parturition) is a time of high oxidative stress, with a potential increase in the requirement for antioxidant nutrients (Masters 2018). Such nutrients are often found in high concentrations in some of the shrubs that are used for shade and shelter (Norman *et al.* 2019). Therefore, in making shelter more attractive, investigations should include the nutritional value of the shelter material and whether the provision of energy, protein or specific nutrients in addition to protection from the wind chill will increase the use of a shelter. Familiarity, through nutritional value and novelty, should be considered as a means of attraction. The ewe faces behavioural decisions on whether to remain with the lamb or to leave to seek a nutritional benefit.

The successful adoption of shelter by ewes at lambing will reduce cold stress but may not be sufficient on its own to take full advantage of potential benefits. For this, the shelter will also need to increase the time that ewes spend at the birth site and encourage interaction between the lamb and ewe. The survival of lambs, particularly multiples, depends on the establishment of a strong ewe–lamb bond which decreases the incidence of separation (Nowak 1996). The development of a strong bond between ewe and lamb is considerably improved if the mother remains on the birth site for a minimum of 6 h after birth (Nowak and Poindron 2006). Merino ewes tend to spend only a short time near the birth site and, consequently, up to 40–50% of twin

lambs can become separated from their mother. Lambs are a partner in this process and in early life (12–24 h) can identify their mother only at close range (0.05 m) (Nowak 1996). When considering shelter, the visual distance between lamb and ewe may be more important than the linear distance (Knight *et al.* 1989), meaning that short or porous shelter would be preferable to high, dense shelter. In line with this suggestion, ewes travelled less in the 24 h after birth when shelter was provided as porous shrubs than when it was 1 m high hessian (Broster *et al.* 2017) and there was 17% less contact between ewes and twin lambs with hessian shelter than with shrub shelter (Broster *et al.* 2010).

Heat

Some studies on the behaviour of sheep offered shade have provided evidence that voluntary shading is not a priority, even at maximum ambient temperatures up to 50°C (Johnson 1987, 1991; Sherwin and Johnson 1990; Johnson and Strack 1992). Conversely, Stafford Smith *et al.* (1985) demonstrated that a respiration index is correlated with shade-seeking behaviour and that this behaviour is further modified by the need for feed and water. The results of Stafford Smith *et al.* (1985) are more consistent with the common site of flocks of sheep resting in the shade of large trees on hot days. To our knowledge, the impact of fleece length on shade-seeking behaviour has not been investigated. It is likely that fleece length would influence the use of shade because it affects radiation heat load and the capacity of sheep to dissipate surface heat. Wool length up to 4 cm provides insulation and reduces radiant heat load (Al-Dawood 2017). However, Merino sheep with a long fleece length (>8 cm) were less able to regulate body temperatures than recently shorn sheep when facing hot but humid conditions (33°C dry bulb temperature; 55% relative humidity; Beatty *et al.* 2008).

Technology to monitor animal behaviour, health, and welfare and the physical environment

Weather stations have been used for many years to collect data that are used to characterise the grazing environment. Such information, sourced from regional meteorological stations, is the basis to the chill index used to alert graziers to the risk of death in lambs and shorn sheep (Nixon-Smith 1972; Donnelly 1984). Technological advancement has meant miniaturisation and reduced unit costs to allow microclimate comparisons within and between sheltered and unsheltered paddocks (Robertson *et al.* 2011; Broster *et al.* 2017). For a useful assessment of shelter, information is also required on the interaction between animals and shelter. Much of the published research on this interaction has been the result of meticulous manual observation and recording (Alexander *et al.* 1979; Pritchard *et al.* 2021). While this is of great value, it is limited by the need for

labour, is restricted to daytime application and observation may influence behaviour. More recently, GPS collars have been used compare the movements of ewes and lambs (Taylor *et al.* 2011; Dobos *et al.* 2015; Broster *et al.* 2017) and contact loggers have been used to compare interactions between ewes and lambs across different types of shelter (Broster *et al.* 2010, 2012b). Now there is information not just on mortality with different weather conditions, but an understanding of the changes in behaviour of the ewes and lambs in response to shelter options. More is possible; for example, accelerometers can provide information on grazing behaviour as well as walking and resting (Giovanetti *et al.* 2017) and intraruminal temperature loggers have been used to monitor drinking events (Vazquez-Diosdado *et al.* 2019).

Technology is now also available to directly monitor the health and welfare of ewes and lambs under experimental conditions using implanted or indwelling devices, such as probes under the skin that extend deep into animal tissues and devices that lodge in the digestive tract. Some of the devices weigh a few grams and have a battery life of 5–10 years (Maloney *et al.* 2019; Lewis Baida *et al.* 2021). Tympanic temperature sensors are being used in cattle to detect fever related to illness (Lewis Baida *et al.* 2021). The deep-core temperature is the gold standard to assess heat stress in homeotherms (Maloney *et al.* 2019). Measurement of skin-surface temperature, subcutaneous temperature or tympanic temperature must be considered carefully because they can be affected by local convection and conduction. Nevertheless, the relationship between the measured internal and external environments, and the interactions of the animal with the physical environment may provide a further indicator of stress and welfare.

Shade, shelter and nutrition

Edible and non-edible shelter

The most common forms of tested shelter are trees, shrubs, grass, vegetative crops, and hessian barriers (Bird *et al.* 1992; Glover *et al.* 2008; Broster *et al.* 2010, 2017). Trees (particularly when mature), rocks and gullies and hessian provide little or no edible dry matter; all other options are edible. Vegetative crops and forages are edible and very palatable, shrubs are usually edible while nutritional value and relative palatability are variable, and mature grass hedges or tussocks are edible but, when mature, have poor nutritional value and are unpalatable. In addition, vegetative crops and forages tend to be uniformly spread over the paddock while shrubs tend to be planted in block or hedgerows. These differences provide opportunities for innovative shelter design; for example, shrubs such as *Atriplex* spp. may provide shelter and are eaten by grazing sheep, other shrubs such as *Rhagodia preissii* provide shelter but are not a preferred diet option (Revell *et al.* 2013),

a shelter with a combination of these shrubs may be attractive to sheep as a food source but also will remain as a functional shelter because of the unpalatable *R. preissii*.

Nutrients from edible shelter

Plants that provide shelter, shade and a feed supply may be more attractive to animals than are shelter and shade with no nutritional value. They may improve the nutrition of the sheep and, at lambing, encourage the ewe to stay in the vicinity of the lamb. While this seems a logical hypothesis, it is not one that has been tested. Of the edible plants, shrubs and trees will provide both shade and shelter while grass hedges and vegetative crops will have some value as shelter but are either too short to provide shade or do not grow during hot seasons of the year. Many herbaceous plants elongate and become taller as they move from vegetative to reproductive phases, this means that shelter value is increasing as nutritional value declines (Norman *et al.* 2021).

Shrubs and trees

Shrubs that are suitable for shelter vary in edible biomass production, nutritive value and palatability. Lefroy *et al.* (1992) reviewed forage trees and shrubs in Australia and concluded that the cultivated species most likely to provide nutrient for livestock include tagasaste (*Chamaecytisus palmensis*) and the tropical species *Leucaena leucocephala*. Robertson and colleagues (Robertson *et al.* 2011; Broster *et al.* 2017) used *Acacia* spp. in a series of studies. *Acacia* spp. usually have a very low digestibility (organic-matter digestibility 27–57%, usually <50%) (Norman *et al.* 2010a; Revell *et al.* 2013) and are unlikely to provide enough ME to contribute significantly to growth and milk production. There are many other shrub options, some with digestibilities high enough to support a moderate level of production and crude protein (CP) well above the requirements for reproducing sheep (Wilson 1977; Lefroy *et al.* 1992; Revell *et al.* 2013). The amount of biomass produced by most shrubs is small compared with pasture production and they will provide a supplement across a range of environments rather than a standalone feed source (Lefroy *et al.* 1992; Ben Salem *et al.* 2010). The value as a supplement may be significant, because these shrubs contain macronutrients, essential micronutrients and secondary compounds that may have both nutritional benefits and nutraceutical properties. Many contain high concentrations of vitamin E and other antioxidants (Pearce *et al.* 2005; Fancote *et al.* 2013; Norman *et al.* 2019), others have been shown to contain compounds that have anthelmintic properties and reduce methane production during rumen fermentation (Kotze *et al.* 2009; Vercoe *et al.* 2009). Mineral composition is also quite different from that of traditional pasture plants (Revell *et al.* 2013). Providing this diversity of nutritional and paranutritional options within a grazing system has been advocated by Provenza and Villalba (2010) as a strategy to enhance

health, intake and preference and by Altieri *et al.* (2015) to provide a more resilient landscape. Evidence that the plants are attractive and provide a novelty factor has been demonstrated, with sheep choosing to make shrubs part of their diet even when there is unlimited high-quality pasture available (Norman *et al.* 2010b).

Trees in Australia do not usually provide a useful supply of nutrients to grazing ruminants. There is often edible biomass, for example, from fallen leaves, fruit, or edible pods but these are usually low in quantity and digestibility. Experiments with grazing sheep in agroforestry systems under *Pinus radiata* have indicated that pine needles have a very low digestibility (~36%) and that fallen pine needles will decrease the quality of pastures growing under the trees (Hawke *et al.* 1993). Conversely, in arid regions of the eastern Mediterranean and Asia, mulberry leaves (*Morus* spp.) are considered a high-value supplement for ruminants (Sánchez 2002).

Herbaceous pasture plants

Grasses and other herbaceous forage plants must be left ungrazed until they are high and dense enough to provide a windbreak at lambing. This may be for most of the growing season and, under drought conditions, there may be insufficient growth to offer shelter. For grasses, this means the tussocks that provide the best shelter value are likely to be rank, have low ME and CP and be high in fibre. For example, tall wheat grass (*Thinopyrum ponticum*) at flowering has a dry-matter digestibility (DMD) of ~46% and CP of 3.4–3.6% (Smith *et al.* 1994). This is well below the requirements for the liveweight maintenance of a non-pregnant adult ewe of approximately 60% DMD and 11% CP (Freer *et al.* 1997). If the plants are managed to prevent flowering, DMD may be 60–85% with CP 15–30%. Other perennial grasses have similar nutritive value; for example, *Phalaris aquatica* (Toowoomba canary grass) and *Lolium perenne* (perennial ryegrass) harvested after anthesis have DMD below 40% (Lam *et al.* 2003). *Bituminaria bituminosa* (tedera) is an untested option; it is an erect plant with a DMD >60% and CP >12% (Oldham *et al.* 2015).

Herbaceous legumes generally offer greater feeding value than do grasses. Norman *et al.* (2021) compared changes in growth and nutritional value during the lifecycle of many of the perennial legume species that are suited to southern Australia. Of the 20 species examined, there was significant variation in biomass production and nutritional value in later stages of plant maturity. Perennial clover species (*Trifolium repens*, *T. pratense* and *T. hybridum*) had the highest nutritional value during senescence (61–66% DMD and 13–14% CP). These clover species struggle to persist in rain-fed mixed farming systems, may lodge when left ungrazed, and may not be sufficiently tall to provide shelter. Sulla (*Hedysarum coronarium*), lucerne (*Medicago sativa*) and sainfoin (*Onobrichis vicifolia*) persist in dry environments and have upright growth habits. Mature, ungrazed stands offer

DMD of 57–61% and CP of 12–16%. Sulla is interesting as it regrows rapidly from the crown in autumn and reaches a significant height before growth is impeded by cold temperatures in winter. If high feeding value results in patchy shelter, there are commercialised herbaceous perennial legumes with very poor feeding value. For example, hairy canary clover (*Dorycnium hirsutum*) produces biomass with very low nutritional value, even in the vegetative phase (DMD < 50%, CP 16%; Norman *et al.* 2021). Although not commercialised, there are several native perennial legumes that may also offer an opportunity as an ‘edible’ shelter, including cullen (*Cullen australasicum*), *Indigofera australis* subsp. *hesperia* and *Glycyrrhiza acanthocarpa* (Bennett *et al.* 2011; Snowball *et al.* 2021).

Vegetative crops

Young crops provide a good supply of highly digestible forage (>80% DMD) with high ME (>12 MJ/kg DM) and CP (>20%) in the early part of the growing season, often when pasture supply is limited (Masters and Thompson 2016). As crops age, their nutritive value declines but still remains adequate for growth and production (de Ruiter *et al.* 2002). The ME and CP content of the crops exceed the minimum requirements for pregnancy and lactation; however, the mineral composition of some crops is imbalanced. A significant proportion of wheat, oat and barley vegetative samples contain less calcium, magnesium and sodium than is required by pregnant sheep and the mineral ratios are consistent with a risk of metabolic disturbances (such as hypocalcaemia and hypomagnesaemia; Dove *et al.* 2016). There is very little known about the impact of grazing cereal crops during lambing on ewe and lamb survival. If animals are removed from crops prior to stem elongation, the yield of harvested grain is usually maintained (Harrison *et al.* 2011). If the crops are grazed by ewes lambing in autumn of winter, there may be insufficient height to provide shelter; alternatively, if grazed by late winter- or spring-lambing ewes, when crops are high enough to provide shelter, a loss of yield may result. Thus, under conditions where cold stress compromises lamb survival, there may be a trade-off between lamb survival and loss of yield at harvest.

Shelter options for grazing sheep

Shelter prevalence

There are very few data on the amount or adequacy of the shelter that currently exists on sheep farms in Australia and New Zealand. Earlier studies from New Zealand, where cold, wet, and windy weather in sheep regions is common, have indicated that in mid-Canterbury only 1% of land was occupied by shelter (Sturrock 1973) and fewer than 23% of farms had one or more shelter belts (Scales 1994). This information is well out of date due to changes in farming

systems in the region. More recently, the majority of sheep farmers who were surveyed identified shelter as important and indicated they were likely to plant shelter in the future (Fisher 2010). The lack of more information is consistent with a low priority for formalised, constructed or planted shelter or shade.

Shelter design and function

Shelter and shade can provide protection from wind, rain, snow and sun and influence both plant and animal health and production. Farmers aiming to provide shelter for livestock also need to consider the consequences for other enterprises and production rotations (e.g. cropping). The enterprise mix will influence the choice of appropriate design and location; there is no single best option.

Selection

Shelter may be planted, built or natural. Natural shelter includes humps, gullies, logs, rocks or natural vegetation. Planted shelter must be appropriate for the soil and environmental conditions and, in most circumstances, indigenous or locally adapted species should be considered first. As the establishment and maintenance of shelter can be expensive, the most economical path will be through the retention and management of natural trees and shrubs (Ritchie 1988). The most common plants and materials used and tested include trees, shrubs, perennial grass tussocks, hessian barriers (or other material such as iron, timber or hay bales), and, to a lesser extent, vegetative crops. Each option has advantages and disadvantages (Table 2). Biological options must have enough vegetative growth to be effective when required (usually during lambing or heat wave conditions). The shelter must be considered within the context of the whole farming system.

Trees have been used in a number of configurations including a single line, a woodlot or block of trees, forest grazing or agroforestry where trees and pasture are grown together for forestry and grazing (Gregory 1995). Shelter and shade from trees may be preferred where timber is potentially a co-product on the farm (Sturrock 1988). Animals may either be excluded or graze within the trees. Trees reduce wind speed by up to 80% close to the row, and continue to provide a significant windbreak at a distance of 5–10 times the height of the trees (Dronen 1988; Bird *et al.* 1992). Trees are a long-term commitment; however, additional benefits may be gained through oil production, biofuels or carbon credits.

Shrubs offer a shelter and shade option; rows of mixed *Acacia* spp. reduced the maximum wind speed in lambing paddocks by 12–53% at a distance 2.5–10 times the height of the rows (Broster *et al.* 2017). Hedgerows of cypress (*Cupressus macrocarpa*; 5–10 m high) reduced average wind speed by 60–70% within 20 m of the hedge

Table 2. Advantages and disadvantages of different shelter option.

Shelter option	Advantages	Disadvantages
Trees	• Protects large area adjacent to the trees • Shelters crops, forage and soils from wind damage and reduces evapotranspiration • Landscape benefits through reduced wind and water erosion and water-table drawdown for salinity management • Provides shade, shelter and habitat to insects, fauna and livestock • Timber and oil production • Carbon-farming opportunity	• Low edible dry matter and green leaves above stock grazing height • Poor nutritive value • Reduced crop and pasture growth adjacent to the tree • Long-term commitment of land (reduced flexibility) • Difficult to locate in cropping landscapes where seeding and harvesting equipment are wide and operated using guidance • Cost and time lags during establishment
Shrubs	• Edible dry matter of low to medium nutritive value but with a high content of some nutrients (e.g. antioxidants and minerals) • Provides both shade and shelter • Correct design will allow visibility between ewe and lamb • Palatable and unpalatable shrub mix to manage vegetation • Landscape benefits through reduced wind and water erosion and water-table drawdown for salinity management • Provides shade, shelter and habitat to insects, fauna and livestock • Low opportunity cost with shrubs that grow on saline, sandy or uncroppable soils • Feed and shelter during seasonal and long-term drought	• Nutritional constraints (e.g. high salt) and secondary compounds (oxalate, saponins and tannins) • Mostly local protection with some benefits to adjacent land • Medium-term commitment of land (reduced flexibility) • High up-front cost of establishment if nursery-raised seedlings are used • Time lags during establishment • Increased management complexity as small-scale equipment may be needed for management of the understorey
Grass (tussocks)	• Good protection from wind adjacent to tussocks • Some edible dry matter • Usually allow visibility between ewe and lamb (may require management) • Easy to remove or relocate	• Tussocks often rank with poor nutritive value • Paddock to be vacated before use to allow tussocks to grow. Results in an economic loss from delayed grazing • Most likely option is to set up as a small, designated lambing paddock • Will offer shelter from wind but no shade • No growth during drought
Vegetative crops and forages	• Very flexible, relatively inexpensive to establish and fits within a mixed farming system • Grain or forage harvest following grazing • High edible dry matter • High nutritive value • May allow visibility between ewe and lamb	• Limited research validation • Loss of harvestable crop yield (plant genotype, growth stage and grazing intensity-dependent) • Nutritional imbalances when grazing monocultures (e.g. minerals in wheat) • Management difficulty, high intakes can quickly reduce shelter value
Hessian, hay bales or man-made structures	• Effective wind break • Flexible construction • Baled straw can be an inexpensive cropping by-product	• Small scale • Often solid, less effective in improving ewe and lamb interaction than are shrubs • Solid structures can create a vortex and cause erosion

(Egan *et al.* 1972). Sheep can also intermingle with the vegetation and find both isolation and protection. Shrubs require a medium- to long-term commitment, although are easier to remove than are large trees and are more suitable for low-rainfall regions where tree growth is slow and perennial grasses and legumes have poor survival. Shrubs such as saltbush and tagasaste have an additional benefit of growing on land that is too saline or sandy for conventional pasture and crop species. Adoption therefore has a lower opportunity cost and the shrubs provide environmental benefits such as reduced salinity and erosion (Masters *et al.* 2005).

Perennial grasses, usually overgrown and mature to form tussocks, have been well investigated and reviewed (Pollard 2006). A 50% reduction in wind speed at lamb height was reported by Cresswell and Thompson (1964); similarly, Lynch and Alexander (1976) reported a reduction of 30–100% at sheep (45 cm) and lamb (10 cm) height in the

lee of tall-grass shelters. While perennial grasses can be very effective, they may fail to grow or regenerate during drought conditions and then provide little shelter (Robertson *et al.* 2011). Perennial legumes have not been investigated to the same degree as have grasses. For both, management will need to be considered, because forage species and accessions within species differ in their nutritional value and growth rates according to planting density, cutting height and cutting frequency (Bell *et al.* 2018; Norman *et al.* 2021).

Vegetative crops are usually grazed during late gestation when crop height is short, and the livestock are removed when stem elongation begins. Longer grazing times with taller crops is an option that would provide both feed and a potential windbreak. In one study, grazing an oat crop that was planted in laneways to allow twin lambs to follow their mother, a non-significant increase in lamb survival was reported compared with a lambing group that grazed pasture (Glover *et al.* 2008). This was during mild winter conditions,

with only a 6–7 cm increase in height of the crop compared with pasture. Anecdotal reports from producers suggest that if the crop is too high, lamb mortality increases due to mismothering, with ewes losing their lambs in the crop, or due to exposure as the ewes lamb along the fence line outside of the shelter provided from the crop. Winter crops will provide no shade benefits during summer and there is no information available on sheep grazing summer crops.

Man-made physical barriers can be highly effective. One metre high hessian rows placed 20 m apart and aligned perpendicular to the prevailing wind significantly reduced wind speed (70–83%) and chill index (5%) over a distance 2–10 times the height of the barrier (Broster *et al.* 2017). Installation can be expensive and time consuming and man-made shelter is better suited to small-scale lambing paddocks. Baled crop stubbles or hay offer another opportunity for shelter and may be less expensive and easier to construct. Many of the options for man-made wind breaks are short and would provide little shade.

Placement

Planning begins with an understanding of the wind across different seasons, with shelter placed to provide protection from the prevailing wind at the time of the highest risk to livestock. Thus, if the prevailing wind is from the west, shelter should be on the western side of the area to be protected (Dronen 1988). This is not always a simple task as climatic data for the time of lambing are not always available and winds that are light or warm will have little impact on survival. This can be addressed by constructing a wind rose to describe the prevailing winds during periods of low temperature and rain (McCaskill and Clark 2007). To minimise wind, shelter should not be planted on ridges as placement there accentuates air flow and may increase wind on the lee side. This is less relevant where slopes are less than 8% (Finch 1988).

There is support for an alternative or complementary approach that is based on an understanding of where ewes choose to lamb without shelter. Alexander *et al.* (1990) observed that Merino ewes tended to seek isolation in preferred lambing areas at higher elevation within a paddock. This was consistent across years. British breed ewes did not show the same tendency and lambed on the edge of the paddock or in association with features such as rocks and trees. The authors acknowledged the confounding factors such as prior history and management of the animals, weather during the studies, paddock size, topographic and other features that make each paddock unique but, even allowing for these, the non-random distribution of birth sites over several years suggested that providing shelter at sites that are naturally selected by the ewes could be effective.

A combination that addresses prevailing wind and weather together with preferred lambing sites may offer the best option.

Design

Shelter value is influenced by the height, density and length of the windbreak (Caborn 1957). On the windward side of a windbreak, wind-speed reductions are measurable upwind for a distance of two to five times the height of the windbreak. On the leeward side, wind-speed reductions occur up to 30 times the height, downwind of the barrier. If the windbreak is very dense, low pressure develops in the leeward side, pulling air downward, thereby creating turbulence and reduced protection (Caborn 1957). A shelter density of 50–80%, with a row spacing of 1–2.5 m for shrubs and 2–6 m for trees has been recommended as the most desirable windbreak density for livestock (Dronen 1988; Finch 1988). This density provides a permeable barrier and more protection; there is a faster recovery in wind speed from a dense shelter belt (Gregory 1995). For shrubs, a complex construction has better shelter effects than does a uniform monoculture (Wu *et al.* 2015). Where shelter is provided in rows, spacing between grass rows of 20–240 m has been effective. If trees or shrubs are used, the protection (and therefore row spacing) can be estimated from the height, or, if the rows are to be combined with alley farming, (Kang *et al.* 1990), machinery width needs to be considered. There is a trade-off between row width and ability to maintain a high-value understorey for production and provision of a diverse diet.

Utilisation

As well as managing shelter to suit livestock, options exist to manage livestock to optimise the potential benefits of shelter or the risks of cold and heat. For example, pregnancy scanning ewes and allocation of twin lambers to shelter is a better strategy than random allocation, particularly if available shelter is limited. Synchronisation of lambing with growth of crop or pasture shelter may improve both shelter and forage supply, and time of shearing may be used to influence shelter use in high-risk environments.

Ancillary effects of shelter and shade

There is a range of secondary benefits from the provision of shelter and shade. Some of these have been identified and quantified, others have not.

Drought and out-of-season feed

Edible shelter and shade have the potential to provide feed during seasonal fluctuations in plant growth and during drought. To be of value, the plants need to be accessible, palatable and be a source of nutrients. Shrubs have long been considered a potential forage source for grazing ruminants, particularly in dry and seasonally variable agricultural zones (Lefroy *et al.* 1992; Le Houérou 1992). Up until the

1990s, only leucaena (*L. leucocephala*) and tagasaste (*C. palmensis*), both species introduced into Australia, were options that had been commercially cultivated (Lefroy *et al.* 1992). Most of Australia's sheep are in the south of the country where only dry and dead forages are available for grazing over summer. Hay or grain supplements are often required at this time. Within this region, tagasaste can partially fill the summer feed gap and is equivalent to or better than production from dry feed and supplements (Oldham *et al.* 1991). Lefroy *et al.* (1992) suggested there are several million hectares of infertile sandy soils in southern Australia where tagasaste could fill a niche.

More recently there has been renewed interest in other shrubs for forage and shelter. Revell *et al.* (2013) measured the nutritive value of 39 Australian native shrub species and found that many had digestibility and CP content high enough to support growth in young sheep. In subsequent experiments, sheep that grazed shrub rows with annual pasture between rows increased liveweight while animals grazing dry pasture alone maintained only weight (Revell *et al.* 2013). The authors suggested that shrubs provide a timely and predictable supply of limiting nutrients with additional benefits in gut health, shade and shelter and could make a valuable addition to the feed base in low- to medium-rainfall zones in southern Australia. The shrub of most interest is saltbush (*Atriplex* spp). Saltbush is a salt-tolerant and drought-tolerant shrub that has been investigated over many years as a feed source and drought reserve in the rangelands (Wilson 1994). The shrub has come back into focus as an option to revegetate saline land in the agricultural zones (Norman *et al.* 2013). *Atriplex nummularia* has now been through a plant-improvement program to develop a cultivar (Annameka) with improved biomass production, palatability and nutritive value relative to most of the wild varieties (Masters and Norman 2010; Norman *et al.* 2016). The availability of these and other halophytic and drought-tolerant plants (Norman *et al.* 2013) provides an opportunity to use land with a low production potential to provide shelter, shade and drought feed.

Vegetative crops are not a drought feeding option as they would usually fail to grow. However, early growth of winter crops at the end of the annual seasonal drought in Mediterranean environments provides an early season feed supply and allows for deferment of grazing in pasture paddocks (Thomas *et al.* 2015).

Pasture and crop production

Providing shade and reduced wind can improve both the rate of growth and the length of the growing season in crop and pasture production. In cold environments, a reduction in wind speed will result in higher temperatures in sheltered plants, leading to improved growth and reduced risk of frost damage (Bird *et al.* 1992). Lower wind speed also means reduced physical damage such as shaking, leaf

stripping or abrasion from blown soils (Nuberg 1998). Shaking reduces cell expansion and growth and has been associated with a 20% reduction in leaf extension area in tall fescue (*Festuca arundinacea*) (Grace 1988). Over the longer term, a reduction in wind speed may reduce evapotranspiration and slow the depletion of soil moisture (Ritchie 1988). This is not a consistent observation as higher wind speed can also lower soil temperature and reduce soil moisture loss (Gregory 1995). Short or prostrate plants are more tolerant of wind damage than are tall erect plants. The extent of the benefit will be dependent on the reduction in wind and, hence, the height and proximity of the shelter. Trees provide wind protection over a larger area than do shrubs, grasses or artificial barriers, with protection up to 10 times the height of the trees. Only in plants that are very close to trees will growth be reduced due to shading and competition for soil moisture (Radcliffe 1983). Increased pasture and animal production have been reported in small, sheltered plots, possibly by changing the water balance within the plants. The benefits were most apparent at high stocking rates (Lynch and Donnelly 1980). Shelter improved the regrowth of pasture grasses in Scotland (Russell and Grace 1979) and improved the growth and, in some studies, the nutritive value of pasture plants in New Zealand, Netherlands, Hungary, USA and Canada (Radcliffe 1983).

Many of the experiments describing the advantages of windbreaks on crop and pasture production have been conducted in cold climates; the benefits in warmer rain-fed cropping systems common in Australia are more variable (Baker *et al.* 2018). There is consistently a loss of production in the competition zone close to the shelter and this is sometimes but not always exceeded by gains in the sheltered zone. In the 12 Australian studies summarised by Baker *et al.* (2018), all showed a net increase in production under some but not all conditions, often in dry years in areas exposed to high winds. The authors suggested that windbreaks may increase in importance with expected hotter and drier climates with more extreme weather events.

Shelterbelts in close proximity to pastures and crops may also act as reservoirs for predatory insects. Tsitsilas *et al.* (2006) reported low numbers of pest mites and lucerne fleas in shelterbelts and increased numbers of predatory mites and spiders both within shelterbelts and nearby pastures. Although others have reported similar benefits (Landis *et al.* 2000), this is not always the case (Dix and Leatherman 1988).

In hot environments, shading and windbreaks will reduce the deleterious effects of higher plant temperature. Shelter may also reduce evapotranspiration and improve the conservation of soil moisture (Gregory 1995).

Landscape health

The role of shelter plants in protecting soils and vegetation has been well summarised in previous reviews (Ritchie 1988;

Bird *et al.* 1992; Lefroy *et al.* 1992; Nuberg 1998). The review by Bird *et al.* (1992) is now 30 years old, but the problems are now worse and solutions more urgent. The authors cited vast areas in Australia's south where wind and water erosion, dryland salinity, soil acidification, loss of soil structure and nutrients and water repellence are causing landscape degradation, costing millions of dollars. Others have described the disproportionate losses of nitrogen, phosphorus and organic carbon in eroded dust (Leys *et al.* 1997). The loss of trees contributes to much of this degradation and means reduced soil organic matter and water-holding capacity and loss of topsoil. Similar problems have been described in the USA, Canada, Russia, China, and Europe (Bird *et al.* 1992). Trees within paddocks also have the potential to provide habitat to complement remnant vegetation in the agricultural landscape and provide transport corridors for birds (Law *et al.* 2000; Fischer and Lindenmayer 2002). The loss of trees is not the only problem, with the loss of shrub and other natural vegetation from arid and semi-arid environments causing similar land and habitat degradation.

Shelterbelts of trees, shrubs, and perennial grass tussocks will not alleviate landscape degradation but can contribute through the modification of wind and water movement. Even single rows or small belts of tall trees can have an impact on wind speed over a large area. Shrubs and grasses will modify wind speed over a much smaller area but, used strategically, can significantly influence water movement and accumulation. Shrubs, trees, and deep-rooted perennial plants have been used to reduce inundation, waterlogging, and flooding, while salt-tolerant shrubs are grown to provide livestock feed and shelter on the saline valley floors (Masters *et al.* 2006).

The configuration of tree and shrub revegetation that is best for landscape health may not also be the best for livestock. For example, the inclusion of livestock may decrease the effectiveness of the trees in reducing wind by reducing height and density of the plants (Bird *et al.* 2007); nevertheless, the publications that have focussed on shelter and shade indicate collectively that there can be benefits for both livestock and the landscape. This should be a consideration in siting and designing shelter and shade.

Practicality and economics of shelter and shade establishment and use

Disadvantages of shelter

The low adoption of shade and shelter indicates that for many sheep producers, active establishment is not a priority. This would indicate that they are not convinced that the advantages outweigh the perceived disadvantages. Disadvantages include the following:

- Reduced flexibility of land use
 - The removal of some land from crop rotation with the establishment of trees, shrubs, and even grass hedges
 - The medium–long-term commitment in the establishment of trees, shrubs and structures can reduce production and management flexibility
 - Physical interference in the use of some of the large semi-autonomous cropping machinery that is used during harvesting, seeding, and spraying
- Loss of productive land
 - The loss of land for crops or pasture. Potential reduced stocking rate
 - The lower productivity of crops and pasture close to shelter due to shading and competition
 - Increase weed-control costs (Walpole 1999)
- Cost of establishment and maintenance
 - Costs for plants, seeds, seedlings, fertiliser, planting, management and fencing
 - Additional management time required to ensure appropriate grazing and livestock nutrition
 - Cost of pruning and thinning
- Soil attributes
 - Compaction and pugging around shelter and shade
 - Nutrients relocated to shade and shelter
- Uncertainty regarding use by sheep and production benefits
 - Over-utilisation of lambing paddocks, with wind speed reducing shelter by over-stocking the density of ewes can lead to reduced lamb survival (Robertson *et al.* 2012)

Parris *et al.* (2011) acknowledged the risk that corridor plantings can cost producers in loss of income, on-going management costs, potential opportunity costs of unforeseen future opportunities, and significant upfront transaction costs. While many of these disadvantages can be addressed, a medium–long-term farm plan is required, together with a detailed land capability assessment.

Economic evaluation of shelter and shade

Estimating the value of shade or shelter to an enterprise or a whole-farm business is particularly challenging, including factors such as existing (remnant) vegetation, its location in relation to pastures, aspect, slope and suitability to provide the primary livestock needs for protection from either the wind or sun, or both. In certain locations, the economic benefit for animal production from shade or shelter will be upscaled when the structure provides adequate protection at the time when reproduction is sensitive to climatic perturbation.

McEachern and Sackett (2008) used a gross-margin analysis to conclude that the cost of the establishment of shelter for the purpose of improving lamb survival was not economically justified. Subsequently, Young *et al.* (2014)

concluded that the McEachern and Sackett (2008) analysis had assumed large costs for establishment, fencing, grazing deferment and stocking rate, with only modest benefits in lamb survival. Young et al. (2014) then published a more comprehensive whole-farm economic analysis that focussed on the use of tall-wheatgrass shelter to improve the survival of winter- and spring-born twins in south-eastern Australia. The study evaluated different flock structure, twinning, mortality, and commodity-price scenarios (2500 scenarios were tested in total). Under all scenarios, the provision of shelter was profitable, ranging from a few cents per ewe to A\$11.35 per ewe. The most beneficial scenario was associated with a 35% increase in the weaning rate, while the least beneficial was associated with a 1% increase. The modelling did not allow for any improvement in the survival of single lambs or for any non-livestock benefits. The limitation of this analysis was that it was based on one location and therefore one climate/production dataset.

Other studies have assessed the economic benefits of trees and shrubs for out-of-season forage supply. A plantation of tagasaste shrubs was shown to be a profitable substitute for hand-feeding grain in autumn (Oldham et al. 1991; Lefroy et al. 1992) and farming 10% of a typical central-wheatbelt crop and livestock property in Western Australia with shrubs has been predicted to increase whole-farm profit by approximately 24% (Monjardino et al. 2010). In Tasmania, measured increases in pasture production were used to model the gross margins from the integration of trees and livestock. Gross margins from a 2-row tree belt that accounted for values for amenity, carbon sequestration, timber production as well as pasture and livestock were 41–74 A\$/ha.year more than for a pasture-only scenario (Mendham 2018). Similarly, Walpole (1999) reported that gross margin of pasture outputs was maximised when the proportion of the tree area across a farm was 34%. The same group (Walpole et al. 1999) surveyed north-eastern Victorian and New South Wales (NSW) Murray-region landholders to report the on-farm benefits and costs associated with remnant native vegetation. Most NSW landholders (>60% of survey participants) were sheep, cattle and cropping enterprises. For these landholders, the most important economic benefits of remnant native vegetation were stock and crop shelter, while for all respondents, the cost of weed control was the major economic cost. Among the NSW participants, 84% used the remnant vegetation for shade and shelter and 95% identified the benefits of remnant native vegetation included its aesthetics and 73% identified benefits for increased stock production. From this work, Walpole et al. (1999) estimated that the value of shade and shelter was A\$4.41/ha and A\$9.54/ha for reduced land degradation. Other benefits, such as cleaner water, nutrient cycling and habitat for pest-controlling predators were identified but by fewer respondents (49%, 42% and 61% respectively).

There may be additional benefits when shade and shelter are provided by native vegetation. Native vegetation

underpins many social and economic aspects of Australian society and has a crucial role in sustaining ecosystem function and processes (Council of Australian Governments 2012). Costanza et al. (2014) estimated that biodiversity contributed more than US\$125 trillion of value to the global economy in 2011. However, trends in Australia have shown a substantial reduction in native vegetation. For example, the Victorian 2018 State of the Environment Report (Commissioner for Environmental Sustainability Victoria 2018) claimed a total loss of 567 913 ha of landscapes supporting native vegetation between 1990 and 2015, primarily from urban development, timber production and agriculture. Some of these losses were native scattered trees (149 063 ha, approximately 27% lost) and native shrubs (48 642 ha, approximately 29% lost).

The value of natural forms of shade and shelter will also have amenity values and may improve real estate value by between 10% and 30% (Oates and Clarke 1987). Those authors were unable to place a value on shade, although recognising its many benefits. The manual *Towards sustainable grazing* (Mason et al. 2003) claims the most successful grazing businesses deliberately incorporate management of environmental issues into their core operation, such as increasing biodiversity using native species and aiming for a minimum 10–15% of the property with conservation as the priority, or including at least one patch of remnant vegetation ≥ 10 ha in size. However, they also recommend that biodiversity initiatives should be focused on the less productive parts of the farm (Mason et al. 2003), which is likely to place quality shelter in places away from quality pastures.

Recommendations for the design and use of shelter and shade – current knowledge base

The assessment of available information indicated that shelter and shade will improve welfare and profit when they are used in locations with a high risk of cold when ewes are lambing or after shearing, or high risk of heat during joining, gestation, and the perinatal period. Further, changing microclimate can lead to other productivity and environmental benefits.

Options that are based on the best current knowledge include the following:

- Shrub belts, preferably using shrubs with a nutritional value to complement pasture and of varying height and palatability. Such belts need to be designed to obstruct the most lethal prevailing wind and consideration should be made to growing them in soils that are unsuitable for crops or at risk of erosion and salinity. Optimum planting would include locally adapted or indigenous species. These shrub belts will also provide shade and a feed supply during dry seasons and droughts.

- Trees, even as a single row, will reduce wind chill, provide shade, and deliver multiple crop and environmental benefits over a paddock or even a landscape scale. To maximise the ancillary benefits, locations should be selected that maximise wind interception.
- Perennial grass hedges and possibly herbaceous legumes are highly effective in improving lamb survival in cold environments. They provide limited feeding value or shade and have traditionally been used on a small scale. This small scale reduces the potential for landscape benefits on erosion and salinity. Grass hedges may be best suited for designated lambing paddocks, particularly those paddocks that are used with twin-lambing ewes.
- Vegetative crops have the theoretical capacity to meet many of the requirements for shelter but not shade. They may be useful and allow flexibility but are so far unproven. Establishment is linked to rainfall and sowing time; producers will need to manage lambing time to match predicted growth patterns.

General principles that apply to all types of shelter include the following:

- A shelter density of 50–80% or a height that will allow some permeability and visual contact between animals.
- That they be located on the side of the paddock perpendicular to the prevailing wind during lambing and away from ridges, particularly in steep terrain.

Extension material on the benefits and costs of shelter and shade will need to include region-specific information on ancillary benefits in plant production and landscape health.

Knowledge gaps – research recommendations

Despite many experiments demonstrating multiple benefits to the planting of shelter and shade for sheep, the practice has been poorly adopted. There are many knowledge gaps, and it is very likely that these are associated with uncertainty and the low adoption.

The following gaps are apparent from this review:

- There is uncertainty around the lamb-survival responses in different regions and climatic conditions. This could be addressed by extending the analysis of [Broster *et al.* \(2012a\)](#) or [McCaskill and Clark \(2007\)](#) to cover all of the major sheep growing climatic zones and lambing strategies. Such information would provide confidence around the potential benefits for decision making.
- Many of the experiments on cold stress have been in small paddocks, leaving a gap in our understanding of the voluntary use of different types of shelter by sheep in large paddocks. This should be combined with

measurement of the behaviour of ewes and lambs and their interaction within shelter. This knowledge is required for both the justification and design of shelter in commercial-sized flocks.

- As part of assessing the use of shelter in a commercial setting, the potential to attract ewes to shelter through the provision of macro and micronutrients and nutraceuticals, should be explored. Such studies should incorporate areas of palatable and unpalatable species.
- Alternative shelter options such as vegetative crops and perennial pasture plants offer options that can be utilised within a routine crop and pasture rotation program and should be investigated.
- The consequences of heat stress on a full production cycle are not well understood. Short-term heat events can alter fertilisation, implantation (returns to service), fetal development and lamb survival. These effects will be insidious and easily missed over an annual production cycle.
- Technology can and should be used within experimental studies to define changes in physiology, stress, and behaviour as the basis for change in production and welfare.
- Collection and compilation of information to allow system benefits (production and ecosystem function) to be modelled and valued across farms and landscapes.

References

- Abe Y, Fujiwara Y, Takahashi H, Matsumura Y, Sawada T, Jiang S, Nakaki R, Uchida A, Nagao N, Naito M, Kajimura S, Kimura H, Osborne TF, Aburatani H, Kodama T, Inagaki T, Sakai J (2018) Histone demethylase JMJD1A coordinates acute and chronic adaptation to cold stress via thermogenic phospho-switch. *Nature Communications* 9, 1566. doi:10.1038/s41467-018-03868-8
- Agricultural Research Council (1980) 'The nutrient requirements of ruminant livestock.' (Commonwealth Agricultural Bureaux: Slough, UK)
- Al-Dawood A (2017) Towards heat stress management in small ruminants – a review. *Annals of Animal Science* 17, 59–88. doi:10.1515/aoas-2016-0068
- Alexander G (1962) Temperature regulation in the new-born lamb. V. summit metabolism. *Australian Journal of Agricultural Research* 13, 100–121. doi:10.1071/AR9620100
- Alexander G, Lynch JJ, Mottershead BE (1979) Use of shelter and selection of lambing sites by shorn and unshorn ewes in paddocks with closely or widely spaced shelters. *Applied Animal Ethology* 5, 51–69. doi:10.1016/0304-3762(79)90007-5
- Alexander G, Lynch JJ, Mottershead BE, Donnelly JB (1980) Reduction in lamb mortality by means of grass wind-breaks: results of a five-year study. *Proceedings of the Australian Society of Animal Production* 13, 329–332.
- Alexander G, Stevens D, Bradley LR (1990) Distribution of field birth-sites of lambing ewes. *Australian Journal of Experimental Agriculture* 30, 759–767. doi:10.1071/EA9900759
- Altieri MA, Nicholls CI, Henao A, Lana MA (2015) Agroecology and the design of climate change-resilient farming systems. *Agronomy for Sustainable Development* 35, 869–890. doi:10.1007/s13593-015-0285-2
- Andrews A (1997) Pregnancy toxemia in the ewe. *In Practice* 19, 306–314. doi:10.1136/inpract.19.6.306
- Animal Health Australia (AHA) (2016) 'Australian animal welfare standards and guidelines – sheep.' (Animal Health Australia (AHA): Canberra, ACT, Australia)

- Arnold GW, Morgan PD (1975) Behaviour of the ewe and lamb at lambing and its relationship to lamb mortality. *Applied Animal Ethology* 2, 25–46. doi:10.1016/0304-3762(75)90063-2
- Baker TP, Moroni MT, Mendham DS, Smith R, Hunt MA (2018) Impacts of windbreak shelter on crop and livestock production. *Crop & Pasture Science* 69, 785–796. doi:10.1071/CP17242
- Barnes A, Beatty D, Taylor E, Stockman C, Maloney S, McCarthy M (2004) 'Physiology of heat stress in cattle and sheep.' (Meat and Livestock Australia: Sydney, NSW, Australia)
- Beatty DT, Barnes A, Fleming PA, Taylor E, Maloney SK (2008) The effect of fleece on core and rumen temperature in sheep. *Journal of Thermal Biology* 33, 437–443. doi:10.1016/j.jtherbio.2008.07.002
- Bell LW, Moore AD, Thomas DT (2018) Integrating diverse forage sources reduces feed gaps on mixed crop-livestock farms. *Animal* 12, 1967–1980. doi:10.1017/S1751731117003196
- Bennett RG, Ryan MH, Colmer TD, Real D (2011) Prioritisation of novel pasture species for use in water-limited agriculture: a case study of Cullen in the Western Australian wheatbelt. *Genetic Resources and Crop Evolution* 58, 83–100. doi:10.1007/s10722-010-9567-3
- Ben Salem H, Norman HC, Nefzaoui A, Mayberry DE, Pearce KL, Revell DK (2010) Potential use of oldman saltbush (*Atriplex nummularia* Lindl.) in sheep and goat feeding. *Small Ruminant Research* 91, 13–28. doi:10.1016/j.smallrumres.2009.10.017
- Bird PR, Bicknell D, Bulman PA, Burke SJA, Leys JF, Parker JN, Van Der Sommen FJ, Voller P (1992) The role of shelter in Australia for protecting soils, plants and livestock. *Agroforestry Systems* 20, 59–86. doi:10.1007/BF00055305
- Bird PR, Jackson TT, Kearney GA, Roache A (2007) Effects of windbreak structure on shelter characteristics. *Australian Journal of Experimental Agriculture* 47, 727–737. doi:10.1071/EA06086
- Broster JC, Dehaan RL, Swain DL, Friend MA (2010) Ewe and lamb contact at lambing is influenced by both shelter type and birth number. *Animal* 4, 796–803. doi:10.1017/S1751731110000030
- Broster JC, Robertson SM, Dehaan RL, King BJ, Friend MA (2012a) Evaluating seasonal risk and the potential for windspeed reductions to reduce chill index at six locations using GrassGro. *Animal Production Science* 52, 921–928. doi:10.1071/AN12066
- Broster JC, Rathbone DP, Robertson SM, King BJ, Friend MA (2012b) Ewe movement and ewe–lamb contact levels in shelter are greater at higher stocking rates. *Animal Production Science* 52, 502–506. doi:10.1071/AN11264
- Broster JC, Dehaan RL, Swain DL, Robertson SM, King BJ, Friend MA (2017) Shelter type and birth number influence the birth and death sites of lambs and ewe movement around lambing time. *Journal of Animal Science* 95, 81–90. doi:10.2527/jas.2016.0692
- Brown GD (1971) Thermal status of sheep at pasture in western New South Wales. *Australian Journal of Agricultural Research* 22, 797–808. doi:10.1071/AR9710797
- Buckman PG (1982) A survey of post-shearing losses following summer rain in south Western Australia. In 'Losses of sheep after shearing due to adverse weather, Darkan, Western Australia'. (Eds SK Baker, HM Chapman, IW Williams) pp. 11–19. (Australian Society of Animal Production (WA Branch): Perth, WA, Australia)
- Caborn JM (1957) 'Shelterbelts and microclimate.' Forestry Commission Bulletin No 29. (H.M. Stationery Office: Edinburgh, Scotland)
- Celi P, Bush R (2010) Pregnancy, lambing and survival. In 'International sheep and wool handbook'. (Ed. DJ Cottle) pp. 223–257. (Nottingham University Press: Nottingham, UK)
- Chauhan SS, Celi P, Fahri FT, Leury BJ, Dunshea FR (2014a) Dietary antioxidants at supranutritional doses modulate skeletal muscle heat shock protein and inflammatory gene expression in sheep exposed to heat stress. *Journal of Animal Science* 92, 4897–4908. doi:10.2527/jas.2014-8047
- Chauhan SS, Celi P, Leury BJ, Clarke IJ, Dunshea FR (2014b) Dietary antioxidants at supranutritional doses improve oxidative status and reduce the negative effects of heat stress in sheep. *Journal of Animal Science* 92, 3364–3374. doi:10.2527/jas.2014-7714
- Commissioner for Environmental Sustainability Victoria (2018) Victorian state of the environment report 2018. Commissioner for Environmental Sustainability Victoria, State Government of Victoria.
- Costanza R, de Groot R, Sutton P, van der Ploeg S, Anderson SJ, Kubiszewski I, Farber S, Turner RK (2014) Changes in the global value of ecosystem services. *Global Environmental Change* 26, 152–158. doi:10.1016/j.gloenvcha.2014.04.002
- Council of Australian Governments (2012) 'Australia's native vegetation framework.' (Council of Australian Governments: Canberra, ACT, Australia)
- Cresswell E, Thompson W (1964) An introductory study of air flow and temperature at grazing-sheep heights. *Empire Journal of Experimental Agriculture* 32, 131–135.
- Crocker K, Curtis K, Speijers J (2009) 'Times of lambing in Australian flocks – 2005 to 2007.' (Department of Agriculture and Food: Perth, WA, Australia)
- Dennis SM (1974) Perinatal lamb mortality in Western Australia: 1. general procedures and results. *Australian Veterinary Journal* 50, 443–449. doi:10.1111/j.1751-0813.1974.tb06867.x
- de Ruiter JM, Hanson R, Hay AS, Armstrong KW, Harrison-Kirk RD (2002) Whole-crop cereals for grazing and silage: balancing quality and quantity. *Proceedings of the New Zealand Grassland Association* 64, 181–189. doi:10.33584/jnzg.2002.64.2456
- Dix ME, Leatherman D (1988) Insect management in windbreaks. *Agriculture, Ecosystems & Environment* 22–23, 513–537. doi:10.1016/0167-8809(88)90043-6
- Dobos RC, Taylor DB, Trotter MG, McCorkell BE, Schneider DA, Hinch GN (2015) Characterising activities of free-ranging Merino ewes before, during and after lambing from GNSS data. *Small Ruminant Research* 131, 12–16. doi:10.1016/j.smallrumres.2015.06.017
- Doney JM, Gunn RG, Griffiths JG (1973) The effect of pre-mating stress on the onset of oestrus and on ovulation rate in Scottish blackface ewes. *Journal of Reproduction and Fertility* 35, 381–384. doi:10.1530/jrf.0.0350381
- Donnelly JR (1984) The productivity of breeding ewes grazing on lucerne or grass and clover pastures on the tablelands of southern Australia. III. Lamb mortality and weaning percentage. *Australian Journal of Agricultural Research* 35, 709–721. doi:10.1071/AR9840709
- Donnelly JR, Moore AD, Freer M (1997) GRAZPLAN: decision support systems for Australian grazing enterprises – I. Overview of the GRAZPLAN project, and a description of the MetAccess and LambAlive DSS. *Agricultural Systems* 54, 57–76. doi:10.1016/S0308-521X(96)00046-7
- Dove H, Masters DG, Thompson AN (2016) New perspectives on the mineral nutrition of livestock grazing cereal and canola crops. *Animal Production Science* 56, 1350–1360. doi:10.1071/AN15264
- Dronen SI (1988) Layout and design criteria for livestock windbreaks. *Agriculture, Ecosystems & Environment* 22–23, 231–240. doi:10.1016/0167-8809(88)90022-9
- Dumont B, Boissy A (2000) Grazing behaviour of sheep in a situation of conflict between feeding and social motivations. *Behavioural Processes* 49, 131–138. doi:10.1016/S0376-6357(00)00082-6
- Dumont B, Dutrion A, Petit M (1998) How readily will sheep walk for a preferred forage? *Journal of Animal Science* 76, 965–971. doi:10.2527/1998.764965x
- Dutt RH (1964) Detrimental effects of high ambient temperature on fertility and early embryo survival in sheep. *International Journal of Biometeorology* 8, 47–56. doi:10.1007/BF02186927
- Dutt RH, Ellington EF, Carlton WW (1959) Fertilization rate and early embryo survival in sheared and unsheared ewes following exposure to elevated air temperature. *Journal of Animal Science* 18, 1308–1318. doi:10.2527/jas1959.1841308x
- Dwyer CM (2008) The welfare of the neonatal lamb. *Small Ruminant Research* 76, 31–41. doi:10.1016/j.smallrumres.2007.12.011
- Eales FA, Small J (1980) Determinants of heat production in newborn lambs. *International Journal of Biometeorology* 24, 157–166. doi:10.1007/BF02253804
- Egan JK, McLaughlin JW, Thompson RL, McIntyre JS (1972) The importance of shelter in reducing neonatal lamb deaths. *Australian Journal of Experimental Agriculture* 12, 470–472. doi:10.1071/EA9720470
- Egan JK, Thompson RL, McIntyre JS (1976) An assessment of overgrown *Phalaris tuberosa* as shelter for newborn lambs. *Proceedings of the Australian Society of Animal Production* 11, 157–160.
- Elvidge DG, Coop IE (1974) Effect of shearing on feed requirements of sheep. *New Zealand Journal of Experimental Agriculture* 2, 397–402. doi:10.1080/03015521.1974.10427703

- Fancote CR, Vercoe PE, Pearce KL, Williams IH, Norman HC (2013) Backgrounding lambs on saltbush provides an effective source of Vitamin E that can prevent Vitamin E deficiency and reduce the incidence of subclinical nutritional myopathy during summer and autumn. *Animal Production Science* **53**, 247–255. doi:10.1071/AN12039
- Finch SJ (1988) 12. Field windbreaks: design criteria. *Agriculture, Ecosystems & Environment* **22–23**, 215–228. doi:10.1016/0167-8809(88)90020-5
- Fischer J, Lindenmayer DB (2002) The conservation value of paddock trees for birds in a variegated landscape in southern New South Wales. 2. Paddock trees as stepping stones. *Biodiversity & Conservation* **11**, 833–849. doi:10.1023/A:1015318328007
- Fisher MW (2010) Shelter and welfare of pastoral animals in New Zealand. *New Zealand Journal of Agricultural Research* **50**, 347–359. doi:10.1080/00288230709510303
- Freer M, Moore AD, Donnelly JR (1997) GRAZPLAN: decision support systems for Australian grazing enterprises – II. The animal biology model for feed intake, production and reproduction and the GrazFeed DSS. *Agricultural Systems* **54**, 77–126. doi:10.1016/S0308-521X(96)00045-5
- Freer M, Dove H, Nolan JV (Eds) (2007) 'Nutrient requirements of domesticated ruminants.' (CSIRO Publishing: Melbourne, Vic., Australia)
- Friend MA, Bhanugopan MS, McGrath SR, Edwards JH, Hancock S, Loudon K, Miller D, McGilchrist P, Refshauge G, Robertson SM, Thompson AN, Masters DG (2020) Do calcium and magnesium deficiencies in reproducing ewes contribute to high lamb mortality? *Animal Production Science* **60**, 733–751. doi:10.1071/AN17588
- Geytenbeek PE (1962) A survey of post shearing losses. *Proceedings of the Australian Society of Animal Production* **4**, 185–186.
- Giovanetti V, Decandia M, Molle G, Acciaro M, Mameli M, Cabiddu A, Cossu R, Serra MG, Manca C, Rassu SPG, Dimauro C (2017) Automatic classification system for grazing, ruminating and resting behaviour of dairy sheep using a tri-axial accelerometer. *Livestock Science* **196**, 42–48. doi:10.1016/j.livsci.2016.12.011
- Glass MH, Jacob RH (1992) Losses of sheep following adverse weather after shearing. *Australian Veterinary Journal* **69**, 142–143. doi:10.1111/j.1751-0813.1992.tb07486.x
- Glover KMM, Milton JTB, Paganoni BL, Martin GB (2008) Survival of twin lambs born in edible shelter is not increased during mild winter weather in Western Australia. *Animal Production in Australia* **27**, 87.
- Grace J (1988) 3. Plant response to wind. *Agriculture, Ecosystems & Environment* **22–23**, 71–88. doi:10.1016/0167-8809(88)90008-4
- Gregory NG (1995) The role of shelterbelts in protecting livestock: a review. *New Zealand Journal of Agricultural Research* **38**, 423–450. doi:10.1080/00288233.1995.9513146
- Hacker R (2010) Extensive grazing systems. In 'International sheep and wool handbook'. (Ed. DJ Cottle) pp. 507–532. (Nottingham University Press: Nottingham, UK)
- Hales JRS (1973) Effects of exposure to hot environments on the regional distribution of blood flow and on cardiorespiratory function in sheep. *Pflügers Archive* **344**, 133–148. doi:10.1007/BF00586547
- Harrison MT, Evans JR, Dove H, Moore AD (2011) Dual-purpose cereals: can the relative influences of management and environment on crop recovery and grain yield be dissected? *Crop & Pasture Science* **62**, 930–946. doi:10.1071/CP11066
- Hawke MF, Dodd MB (2003) Livestock shelter from trees – a review. *NZGA: Research and Practice Series* **10**, 65–72. doi:10.33584/rps.10.2003.2985
- Hawke MF, Rattray PV, Percival NS (1993) Liveweight changes of sheep grazing a range of herbage allowances under *Pinus radiata* agroforestry regimes. *Agroforestry Systems* **23**, 11–21. doi:10.1007/BF00704848
- Hinch GN, Brien F (2014) Lamb survival in Australian flocks: a review. *Animal Production Science* **54**, 656–666. doi:10.1071/AN13236
- Hopkins PS, Nolan CJ, Pepper PM (1980) The effects of heat stress on the development of the foetal lamb. *Australian Journal of Agricultural Research* **31**, 763–771. doi:10.1071/AR9800763
- Hutchinson KJ (1966) Bodyweight criteria and post shearing deaths. *Proceedings of the Australian Society of Animal Production* **6**, 229–233.
- Hutchinson JCD (1968) Deaths of sheep after shearing. *Australian Journal of Experimental Agriculture* **8**, 393–400. doi:10.1071/EA9680393
- Johnson KG (1987) Shading behaviour of sheep: preliminary studies of its relation to thermoregulation, feed and water intakes, and metabolic rates. *Australian Journal of Agricultural Research* **38**, 587–596. doi:10.1071/AR9870587
- Johnson KG (1991) Body temperatures and respiratory rates of free-ranging Merino sheep in and out of shade during summer. *Australian Journal of Agricultural Research* **42**, 1347–1357. doi:10.1071/AR9911347
- Johnson KG, Strack R (1992) Effects of shade use on grazing, drinking, ruminating and postural patterns of Merino sheep. *Australian Journal of Agricultural Research* **43**, 261–264. doi:10.1071/AR9920261
- Kang BT, Reynolds L, Atta-Krah AN (1990) Alley farming. *Advances in Agronomy* **43**, 315–359. doi:10.1016/S0065-2113(08)60481-2
- Kenyon PR, Morris ST, Revell DK, McCutcheon SN (2003) Shearing during pregnancy – review of a policy to increase birthweight and survival of lambs in New Zealand pastoral farming systems. *New Zealand Veterinary Journal* **51**, 200–207. doi:10.1080/00480169.2003.36368
- Kleemann DO, Walker SK (2005) Fertility in South Australian commercial Merino flocks: relationships between reproductive traits and environmental cues. *Theriogenology* **63**, 2416–2433. doi:10.1016/j.theriogenology.2004.09.052
- Knight TW, Wilson LD, Lynch PR, Hockey H-UP (1989) Slope and the choice of birth sites by ewes. *New Zealand Journal of Agricultural Research* **32**, 193–198. doi:10.1080/00288233.1989.10423453
- Kotze AC, O'Grady J, Emms J, Toovey AF, Hughes S, Jessop P, Bennell M, Vercoe PE, Revell DK (2009) Exploring the anthelmintic properties of Australian native shrubs with respect to their potential role in livestock grazing systems. *Parasitology* **136**, 1065–1080. doi:10.1017/S0031182009006386
- Lam TB-T, Iiyama K, Stone BA (2003) Hot alkali-labile linkages in the walls of the forage grass *Phalaris aquatica* and *Lolium perenne* and their relation to in vitro wall digestibility. *Phytochemistry* **64**, 603–607. doi:10.1016/S0031-9422(03)00301-7
- Landis DA, Wratten SD, Gurr GM (2000) Habitat management to conserve natural enemies of arthropod pests in agriculture. *Annual Review of Entomology* **45**, 175–201. doi:10.1146/annurev.ento.45.1.175
- Law BS, Chidel M, Turner G (2000) The use by wildlife of paddock trees in farmland. *Pacific Conservation Biology* **6**, 130–143. doi:10.1071/PC000130
- Lefroy EC, Dann PR, Wildin JH, Wesley-Smith RN, McGowan AA (1992) Trees and shrubs as sources of fodder in Australia. *Agroforestry Systems* **20**, 117–139. doi:10.1007/BF00055307
- Le Houérou HN (1992) The role of saltbushes (*Atriplex* spp.) in arid land rehabilitation in the Mediterranean Basin: a review. *Agroforestry Systems* **18**, 107–148. doi:10.1007/BF00115408
- Lewis Baida BE, Swinbourne AM, Barwick J, Leu ST, van Wettene WHEJ (2021) Technologies for the automated collection of heat stress data in sheep. *Animal Biotelemetry* **9**, 4. doi:10.1186/s40317-020-00225-9
- Leys JF, Butler P, Packer S, New South Wales Department of Land and Water Conservation (1997) 'Wind erosion research at Keith in the upper south east of South Australia: a component of the 'sustainable agriculture for upper south east farms''. (Department of Land and Water Conservation)
- Lowe TE, Gregory NG, Fisher AD, Payne SR (2002) The effects of temperature elevation and water deprivation on lamb physiology, welfare, and meat quality. *Australian Journal of Agricultural Research* **53**, 707–714. doi:10.1071/AR01125
- Lynch JJ, Alexander G (1976) The effect of graminaceous windbreaks on behaviour and lamb mortality among shorn and unshorn merino sheep during lambing. *Applied Animal Ethology* **2**, 305–325. doi:10.1016/0304-3762(76)90065-1
- Lynch JJ, Alexander G (1977) Sheltering behaviour of lambing Merino sheep in relation to grass hedges and artificial windbreaks. *Australian Journal of Agricultural Research* **28**, 691–701. doi:10.1071/AR9770691
- Lynch JJ, Donnelly JB (1980) Changes in pasture and animal production resulting from the use of windbreaks. *Australian Journal of Agricultural Research* **31**, 967–979. doi:10.1071/AR9800967
- Lynch JJ, Mottershead BE, Alexander G (1980) Sheltering behaviour and lamb mortality amongst shorn Merino ewes lambing in paddocks with a restricted area of shelter or no shelter. *Applied Animal Ethology* **6**, 163–174. doi:10.1016/0304-3762(80)90067-X

- Maloney SK, Goh G, Fuller A, Vesterdorf K, Blache D (2019) Amplitude of the circadian rhythm of temperature in homeotherms. *CABI Reviews: Perspectives in Agriculture, Veterinary Science, Nutrition and Natural Resources* **14**, 1–30. doi:10.1079/PAVSNR201914019
- Marai IFM, El-Darawany AA, Fadiel A, Abdel-Hafez MAM (2007) Physiological traits as affected by heat stress in sheep – a review. *Small Ruminant Research* **71**, 1–12. doi:10.1016/j.smallrumres.2006.10.003
- Mason W, Warn LK, Cahill G (2003) 'Towards sustainable grazing: the professional producer's guide.' (Meat and Livestock Australia: Sydney, NSW, Australia)
- Masters DG (2018) Practical implications of mineral and vitamin imbalance in grazing sheep. *Animal Production Science* **58**, 1438–1450. doi:10.1071/AN17761
- Masters D, Norman H (2010) Salt tolerant plants for livestock production. In 'Proceedings of the international conference on soils and groundwater salinization in arid regions', 11–14 January 2010, Muscat, Oman. (Eds M Ahmed, SA Al-Rawahy) (Sultan Qaboos University: Muscat, Oman)
- Masters DG, Thompson AN (2016) Grazing crops: implications for reproducing sheep. *Animal Production Science* **56**, 655–668. doi:10.1071/AN14517
- Masters DG, Norman HC, Barrett-Lennard EG (2005) Agricultural systems for saline soil: the potential role of livestock. *Asian-Australasian Journal of Animal Sciences* **18**, 296–300. doi:10.5713/ajas.2005.296
- Masters D, Edwards N, Silience M, Avery A, Revell D, Friend M, Sanford P, Saul G, Beverly C, Young J (2006) The role of livestock in the management of dryland salinity. *Australian Journal of Experimental Agriculture* **46**, 733–741. doi:10.1071/EA06017
- Mavrogiani VS, Brozos C (2008) Reflections on the causes and the diagnosis of peri-parturient losses of ewes. *Small Ruminant Research* **76**, 77–82. doi:10.1016/j.smallrumres.2007.12.019
- McCaskill M, Clark S (2007) EverGraze 4. Wind roses for 'bad' lambing weather. In '48th annual conference of the grassland society of southern Australia, Echuca, Victoria', p. 103. (Grasslands Society of Southern Australia)
- McCrabb GJ, Bortolussi G (1996) Placental growth and the ability of sheep to thermoregulate in hot environment. *Small Ruminant Research* **20**, 121–127. doi:10.1016/0921-4488(95)00802-0
- McCrabb GJ, McDonald BJ, Hennoste LM (1993) Heat stress during mid-pregnancy in sheep and the consequences for placental and fetal growth. *The Journal of Agricultural Science* **120**, 265–271. doi:10.1017/S0021859600074311
- McEachern S, Sackett D (2008) 'Economic analysis of management options to improve reproductive performance in Merino ewes.' (Holmes, Sackett & Associates: Wagga Wagga, NSW, Australia)
- McLaughlin JW, Egan JK, Poynton WM, Thompson RL (1970) The effect upon neonatal lamb mortality of lambing systems incorporating the use of partial and complete shelter. *Proceedings of the Australian Society of Animal Production* **8**, 337–343.
- McNeill DM, Kelly RW, Williams IH (1997) Insulin sensitivity and fetal growth in ewes maintained in a moderate body condition compared with a lean body condition in late pregnancy. *Australian Journal of Agricultural Research* **48**, 753–758. doi:10.1071/A96133
- McQuillan M, Glanville E, Jacobson C, Sherriff L, Whale A (2021) Unlocking the keys to ewe survival. In 'Proceedings of the Australian veterinary association – sheep, camelid and goat veterinary conference', Wagga Wagga, pp. 48–50. (Australian Veterinary Association: Australia)
- Mellor DJ, Stafford KJ (2004) Animal welfare implications of neonatal mortality and morbidity in farm animals. *The Veterinary Journal* **168**, 118–133. doi:10.1016/j.tvjl.2003.08.004
- Mendham D (2018) 'Modelling the costs and benefits of Agroforestry systems – application of the Imagine bioeconomic model at four case study sites in Tasmania.' (CSIRO)
- Mitchell D, Snelling EP, Hetem RS, Maloney SK, Strauss WM, Fuller A (2018) Revisiting concepts of thermal physiology: predicting responses of mammals to climate change. *Journal of Animal Ecology* **87**, 956–973. doi:10.1111/1365-2656.12818
- Monjardino M, Revell D, Pannell DJ (2010) The potential contribution of forage shrubs to economic returns and environmental management in Australian dryland agricultural systems. *Agricultural Systems* **103**, 187–197. doi:10.1016/j.agsy.2009.12.007
- Morrison SR (1983) Ruminant heat stress: effect on production and means of alleviation. *Journal of Animal Science* **57**, 1594–1600. doi:10.2527/jas1983.5761594x
- Moslemipur F, Golzar-Adabi S (2017) Physiological and growth parameters of fattening lambs after shearing under heat-stress conditions. *Animal Production Science* **57**, 569–575. doi:10.1071/AN15001
- Mottershead BE, Alexander G, Lynch JJ (1982) Sheltering behaviour of shorn and unshorn sheep in mixed or separate flocks. *Applied Animal Ethology* **8**, 127–136. doi:10.1016/0304-3762(82)90138-9
- National Research Council (1981) 'Effect of environment on nutrient requirements of domestic animals.' (National Academy Press: Washington, DC, USA)
- Nixon-Smith WF (1972) 'The forecasting of chill index ratings for newborn lambs and off-shears sheep by use of a cooling factor derived from synoptic data.' (Bureau of Meteorology: Canberra, ACT, Australia)
- Norman HC, Revell DK, Mayberry DE, Rintoul AJ, Wilmot MG, Masters DG (2010a) Comparison of *in vivo* organic matter digestion of native Australian shrubs by sheep to *in vitro* and *in sacco* predictions. *Small Ruminant Research* **91**, 69–80. doi:10.1016/j.smallrumres.2009.11.019
- Norman HC, Wilmot MG, Thomas DT, Barrett-Lennard EG, Masters DG (2010b) Sheep production, plant growth and nutritive value of a saltbush-based pasture system subject to rotational grazing or set stocking. *Small Ruminant Research* **91**, 103–109. doi:10.1016/j.smallrumres.2009.11.022
- Norman HC, Masters DG, Barrett-Lennard EG (2013) Halophytes as forages in saline landscapes: interactions between plant genotype and environment change their feeding value to ruminants. *Environmental and Experimental Botany* **92**, 96–109. doi:10.1016/j.envexpbot.2012.07.003
- Norman HC, Hulm E, Wilmot MG (2016) Improving the feeding value of Old Man Saltbush for saline production systems in Australia. In 'Halophytic and salt tolerant feedstuffs: impacts on nutrition, physiology and reproduction of livestock'. (Eds HM El Shaer, VR Squires) pp. 79–86. (CRC Press: Boca Raton, FL, USA)
- Norman HC, Duncan EG, Masters DG (2019) Halophytic shrubs accumulate minerals associated with antioxidant pathways. *Grass and Forage Science* **74**, 345–355. doi:10.1111/gfs.12434
- Norman HC, Humphries AW, Hulm E, Young P, Hughes SJ, Rowe T, Peck DM, Vercoe PE (2021) Productivity and nutritional value of 20 species of perennial legumes in a low-rainfall Mediterranean-type environment in southern Australia. *Grass and Forage Science* **76**, 134–158. doi:10.1111/gfs.12527
- Nowak R (1996) Neonatal survival: contributions from behavioural studies in sheep. *Applied Animal Behaviour Science* **49**, 61–72. doi:10.1016/0168-1591(95)00668-0
- Nowak R, Poindron P (2006) From birth to colostrum: early steps leading to lamb survival. *Reproduction Nutrition Development* **46**, 431–446. doi:10.1051/rnd:2006023
- Nuberg IK (1998) Effect of shelter on temperate crops: a review to define research for Australian conditions. *Agroforestry Systems* **41**, 3–34. doi:10.1023/A:1006071821948
- Oates N, Clarke B (1987) 'Trees for the back paddock.' (Goddard and Dobson Publishers: Melbourne, Vic., Australia)
- Obst JM, Day HR (1968) The effect of inclement weather on mortality of Merino and Corriedale lambs on Kangaroo Island. *Proceedings of the Australian Society of Animal Production* **8**, 239–242.
- Oldham C, Allen G, Moore P, Mattinson B (1991) Animal production from tagasaste growing in deep sands in a 450 mm winter rainfall zone. *Journal of the Department of Agriculture, Western Australia* **32**, 24–30.
- Oldham CM, Thompson AN, Ferguson MB, Gordon DJ, Kearney GA, Paganoni BL (2011) The birthweight and survival of Merino lambs can be predicted from the profile of liveweight change of their mothers during pregnancy. *Animal Production Science* **51**, 776–783. doi:10.1071/AN10155
- Oldham CM, Wood D, Milton J, Real D, Vercoe P, van Burgel AJ (2015) An animal house study on utilisation of fresh teder (Bituminaria bituminosa var. albomarginata and crassiuscula) by Merino wethers. *Animal Production Science* **55**, 617–624. doi:10.1071/AN13068
- Parris H, Whitten SM, Wyborn C, Hill R, Freudenberger D (2011) An overview of key socio-economic factors, principles and

- guidelines in wildlife 'corridor' planning and implementation. Australian Government Department of Sustainability, Environment, Water, Population and Communities.
- Pearce KL, Masters DG, Smith GM, Jacob RH, Pethick DW (2005) Plasma and tissue α -tocopherol concentrations and meat colour stability in sheep grazing saltbush (*Atriplex* spp.). *Australian Journal of Agricultural Research* **56**, 663–672. doi:10.1071/AR05001
- Phillips C (2016) The welfare risks and impacts of heat stress on sheep shipped from Australia to the Middle East. *The Veterinary Journal* **218**, 78–85. doi:10.1016/j.tvjl.2016.09.011
- Pollard JC (2006) Shelter for lambing sheep in New Zealand: a review. *New Zealand Journal of Agricultural Research* **49**, 395–404. doi:10.1080/00288233.2006.9513730
- Pollard JC, Littlejohn RP (1999) Shelter for lambing in southern New Zealand. II. Sheltering behaviour and effects on productivity. *New Zealand Journal of Agricultural Research* **42**, 171–177. doi:10.1080/00288233.1999.9513367
- Porter WP, Gates DM (1969) Thermodynamic equilibria of animals with environment. *Ecological Monographs* **39**, 227–244. doi:10.2307/1948545
- Pritchard CE, Williams AP, Davies P, Jones D, Smith AR (2021) Spatial behaviour of sheep during the neonatal period: preliminary study on the influence of shelter. *Animal* **15**, 100252. doi:10.1016/j.animal.2021.100252
- Provenza FD, Villalba JJ (2010) The role of natural plant products in modulating the immune system: an adaptable approach for combating disease in grazing animals. *Small Ruminant Research* **89**, 131–139. doi:10.1016/j.smallrumres.2009.12.035
- Radcliffe JE (1983) Grassland responses to shelter – a review. *New Zealand Journal of Experimental Agriculture* **11**, 5–10. doi:10.1080/03015521.1983.10427718
- Refshauge G, Brien FD, Hinch GN, van de Ven R (2016) Neonatal lamb mortality: factors associated with the death of Australian lambs. *Animal Production Science* **56**, 726–735. doi:10.1071/AN15121
- Reid RL (1962) Studies on the carbohydrate metabolism of sheep. XV. The adrenal response to the climatic stresses of cold, wind, and rain. *Australian Journal of Agricultural Research* **13**, 296–306. doi:10.1071/AR9620296
- Revell DK, Norman HC, Vercoe PE, Phillips N, Toovey A, Bickell S, Hulm E, Hughes S, Emms J (2013) Australian perennial shrub species add value to the feed base of grazing livestock in low- to medium-rainfall zones. *Animal Production Science* **53**, 1221–1230. doi:10.1071/AN13238
- Ritchie KA (1988) 24. Shelterbelt plantings in semi-arid areas. *Agriculture, Ecosystems & Environment* **22-23**, 425–440. doi:10.1016/0167-8809(88)90037-0
- Robertson SM, Friend MA, Broster JC, King BJ (2011) Survival of twin lambs is increased with shrub belts. *Animal Production Science* **51**, 925–938. doi:10.1071/AN11006
- Robertson SM, King BJ, Broster JC, Friend MA (2012) The survival of lambs in shelter declines at high stocking intensities. *Animal Production Science* **52**, 497–501. doi:10.1071/AN11261
- Robinson WI, Brown W, Lucas IAM (1973) Effects of shelter, exposure and level of feeding over winter on the productivity of Welsh Mountain ewes and lambs. *Animal Science* **17**, 21–32. doi:10.1017/S0003356100030245
- Russell G, Grace J (1979) The effect of shelter on the yield of grasses in southern Scotland. *Journal of Applied Ecology* **16**, 319–330. doi:10.2307/2402751
- Sánchez MD (2002) World distribution and utilization of Mulberry, potential for animal feeding. In 'Mulberry for animal production (Morus1-L)'. (Food and Agriculture Organisation of the United Nations: Rome, Italy)
- Sawyer GJ (1979) The influence of radiant heat load on reproduction in the Merino ewe. II. The relative effects of heating before and after insemination. *Australian Journal of Agricultural Research* **30**, 1143–1149. doi:10.1071/AR9791143
- Scales GH (1994) Snowfall risk management 1992 snows. A survey of management practices used by Canterbury farmers to minimise the impact of snow. No. MAF technical paper 94/11, Ministry of Agriculture and Fisheries, Wellington, NZ.
- Sherwin CM, Johnson KG (1990) Skin and abdominal temperatures recorded by data loggers attached to Merino sheep voluntarily staying out of shade. *Australian Journal of Agricultural Research* **41**, 781–790. doi:10.1071/AR9900781
- Silanikove N (1987) Impact of shelter in hot Mediterranean climate on feed intake, feed utilization and body fluid distribution in sheep. *Appetite* **9**, 207–215. doi:10.1016/S0195-6663(87)80014-4
- Silanikove N (2000) Effects of heat stress on the welfare of extensively managed domestic ruminants. *Livestock Production Science* **67**, 1–18. doi:10.1016/S0301-6226(00)00162-7
- Slee J, Ryder ML (1967) The effect of cold exposure on wool growth in Scottish Blackface and Merino $\times$ Cheviot Sheep. *The Journal of Agricultural Science* **69**, 449–453. doi:10.1017/S0021859600019146
- Smith JF (1971) The effect of temperature on characteristics of semen of rams. *Australian Journal of Agricultural Research* **22**, 481–490. doi:10.1071/AR9710481
- Smith ID, Bell GH, de Chaneet G (1966) Embryonic mortality in merino ewes exposed to high ambient temperatures. *Australian Veterinary Journal* **42**, 468–470. doi:10.1111/j.1751-0813.1966.tb14478.x
- Smith KF, Lee CK, Borg PT, Flinn PC (1994) Yield, nutritive value, and phenotypic variability of tall wheatgrass grown in a nonsaline environment. *Australian Journal of Experimental Agriculture* **34**, 609–614. doi:10.1071/EA9940609
- Snowball R, Norman HC, D'Antuono MF (2021) Investigation of two native Australian perennial forage legumes for their potential use in agriculture: *Indigofera australis* subsp. *hesperia* and *Glycyrrhiza acanthocarpa*. *Crop & Pasture Science* **72**, 311–323. doi:10.1071/CP20287
- Sonna LA, Fujita J, Gaffin SL, Lilly CM (2002) Invited Review: Effects of heat and cold stress on mammalian gene expression. *Journal of Applied Physiology* **92**, 1725–1742. doi:10.1152/jappphysiol.01143.2001
- Stafford Smith DM, Noble IR, Jones GK (1985) A heat balance model for sheep and its use to predict shade-seeking behaviour in hot conditions. *Journal of Applied Ecology* **22**, 753–774. doi:10.2307/2403227
- Stephenson RGA, Suter GR, Le Feuvre AS (1984) Reduction of the effects of heat stress on lamb birth weight and survival by provision of shade. In 'Reproduction in sheep'. (Eds DR Lindsay, DT Pearce) pp. 223–225. (Cambridge University Press: Cambridge)
- Stevens D, Alexander G, Lynch JJ (1981) Do Merino ewes seek isolation or shelter at lambing? *Applied Animal Ethology* **7**, 149–155. doi:10.1016/0304-3762(81)90093-6
- Stockman CA, Barnes AL, Maloney SK, Taylor E, McCarthy M, Pethick D (2011) Effect of prolonged exposure to continuous heat and humidity similar to long haul live export voyages in Merino wethers. *Animal Production Science* **51**, 135–143. doi:10.1071/AN10120
- Sturrock JW (1973) Surveys of tree and hedge shelter in mid Canterbury: a summary of results. *New Zealand Journal of Experimental Agriculture* **1**, 105–107. doi:10.1080/03015521.1973.10427627
- Sturrock JW (1988) Shelter: its management and promotion. *Agriculture, Ecosystems & Environment* **22-23**, 1–13. doi:10.1016/0167-8809(88)90004-7
- Taylor DB, Schneider DA, Brown WY, Price IR, Trotter MG, Lamb DW, Hinch GN (2011) GPS observation of shelter utilisation by Merino ewes. *Animal Production Science* **51**, 724–737. doi:10.1071/AN11025
- Thomas DT, Moore AD, Norman HC, Revell CK (2015) Small effects of deferment of annual pastures through grazing spring wheat crops in Western Australia can benefit livestock productivity. *Crop & Pasture Science* **66**, 410–417. doi:10.1071/CP14090
- Thwaites CJ (1967) Prolonged heat stress and wool growth in sheep. *International Journal of Biometeorology* **11**, 297–300. doi:10.1007/BF01426652
- Thwaites CJ (1971) Short term heat stress and embryo mortality in the ewe. *Australian Journal of Experimental Agriculture* **11**, 265–267. doi:10.1071/EA9710265
- Tsitsilas A, Stuckey S, Hoffmann AA, Weeks AR, Thomson LJ (2006) Shelterbelts in agricultural landscapes suppress invertebrate pests. *Australian Journal of Experimental Agriculture* **46**, 1379–1388. doi:10.1071/EA05137
- van Wetteer WHEJ, Kind KL, Gatford KL, Swinbourne AM, Leu ST, Hayman PT, Kelly JM, Weaver AC, Kleemann DO, Walker SK (2021) Review of the impact of heat stress on reproductive performance of sheep. *Journal of Animal Science and Biotechnology* **12**, 26. doi:10.1186/s40104-020-00537-z

- Vazquez-Diosdado JA, Miguel-Pacheco GG, Plant B, Dottorini T, Green M, Kaler J (2019) Developing and evaluating threshold-based algorithms to detect drinking behavior in dairy cows using reticulorumen temperature. *Journal of Dairy Science* **102**, 10471–10482. doi:10.3168/jds.2019-16442
- Vercoe PE, Durmic Z, Revell DK (2009) Rumen microbial ecology: helping to change landscape. *Options Méditerranéennes* **85**, 225–236.
- Walpole SC (1999) Assessment of the economic and ecological impacts of remnant vegetation on pasture productivity. *Pacific Conservation Biology* **5**, 28–35. doi:10.1071/PC990028
- Walpole SC, Miles C, Lockwood M (1999) Assessment of the on-farm economic values of remnant native vegetation. In 'AARES 43rd Annual Conference/NZARES 6th Annual Conference', 19–22 January, 1999, Christchurch, New Zealand. (Australian Agricultural and Resource Economics Society) p. 57.
- Watson RH, Alexander G, Cummings IA, MacDonald JW, McLaughlin JW, Rizzoli DJ, Williams D (1968) Reduction of perinatal loss of lambs in winter in Western Victoria by lambing in sheltered individual pens. *Proceedings of the Australian Society of Animal Production* **8**, 243–249.
- Webster AJF, Osuji PO, White F, Ingram JF (1975) The influence of food intake on portal blood flow and heat production in the digestive tract of sheep. *British Journal of Nutrition* **34**, 125–139. doi:10.1017/S0007114575000165
- Wilson AD (1977) The digestibility and voluntary intake of the leaves of trees and shrubs by sheep and goats. *Australian Journal of Agricultural Research* **28**, 501–508. doi:10.1071/AR9770501
- Wilson AD (1994) Halophytic shrubs in semi-arid regions of Australia. In 'Halophytes as a resource for livestock and for rehabilitation of degraded lands'. (Eds VR Squires, AT Ayoub) pp. 101–113. (Kluwer Academic Publishers: Dordrecht, Netherlands)
- Wu X, Zou X, Zhou N, Zhang C, Shi S (2015) Deceleration efficiencies of shrub windbreaks in a wind tunnel. *Aeolian Research* **16**, 11–23. doi:10.1016/j.aeolia.2014.10.004
- Young BA (1981) Cold stress as it affects animal production. *Journal of Animal Science* **52**, 154–163. doi:10.2527/jas1981.521154x
- Young JM, Saul G, Behrendt R, Byrne F, McCaskill M, Kearney GA, Thompson AN (2014) The economic benefits of providing shelter to reduce the mortality of twin lambs in south-western Victoria. *Animal Production Science* **54**, 773–782. doi:10.1071/AN13256

Data availability. All data used in this paper were sourced from previously published information and available as cited.

Conflicts of interest. The authors declare they have no conflicts of interest.

Declaration of funding. This review was funded as part of Meat and Livestock Australia project P.PSH.1316 and prepared in collaboration with project P.PSH.1292. Meat and Livestock Australia approved the paper for publication but was not involved in the analysis or interpretation of information collected or in the writing of the paper.

Author affiliations

^ASchool of Agriculture and Environment, Faculty of Science, The University of Western Australia, 35 Stirling Highway, Crawley, WA 6009, Australia.

^BCSIRO Agriculture and Food, Private Bag 5, Wembley, WA 6913, Australia.

^CCollege of Science, Health, Engineering and Education, Murdoch University, Murdoch WA 6150, Australia.

^DSchool of Human Sciences, Faculty of Science, The University of Western Australia, 35 Stirling Highway, Crawley, WA 6009, Australia.

^ENew South Wales Department of Primary Industries, Cowra Agricultural Research and Advisory Station, Cowra, NSW 2794, Australia.